Pure and Applied Functional Analysis Volume 4, Number 1, 2019, 21–33



DISCRETE TIME PONTRYAGIN PRINCIPLES IN BANACH SPACES

MOHAMMED BACHIR AND JOËL BLOT

ABSTRACT. The aim of this paper is to establish Pontryagin's principles in a dicrete-time infinite-horizon setting when the state variables and the control variables belong to infinite-dimensional Banach spaces. In comparison with previous results on this question, we delete conditions of finiteness of codimension of subspaces. To realize this aim, the main ideas are the introduction of new recursive assumptions and useful consequences of the Baire category theorem and of the Banach isomorphism theorem.

1. INTRODUCTION

The considered infinite-horizon Optimal Control problems are governed by the following discrete-time controlled dynamical system.

(1.1)
$$x_{t+1} = f_t(x_t, u_t), \quad t \in \mathbb{N}$$

where $x_t \in X_t \subset X$, $u_t \in U_t \subset U$ and $f_t : X_t \times U_t \to X_{t+1}$. Here X and U are real Banach spaces; X_t is a nonempty open subset of X and U_t is a nonempty subset of U. As usual, the x_t are called the state variables and the u_t are called the control variables.

From an initial state $\sigma \in X_0$, we denote by $Adm(\sigma)$ the set of the processes $((x_t)_{t\in\mathbb{N}}, (u_t)_{t\in\mathbb{N}}) \in (\prod_{t\in\mathbb{N}} X_t) \times (\prod_{t\in\mathbb{N}} U_t)$ which satisfy (1.1) for all $t\in\mathbb{N}$. The elements of $Adm(\sigma)$ are called the admissible processes.

For all $t \in \mathbb{N}$, we consider the function $\phi_t : X_t \times U_t \to \mathbb{R}$ to define the criteria.We denote by Dom(J) the set of the $((x_t)_{t \in \mathbb{N}}, (u_t)_{t \in \mathbb{N}}) \in (\prod_{t \in \mathbb{N}} X_t) \times (\prod_{t \in \mathbb{N}} U_t)$ such that the series $\sum_{t=0}^{+\infty} \phi_t(x_t, u_t)$ is convergent in \mathbb{R} . We define the nonlinear functional $J : Dom(J) \to \mathbb{R}$ by setting

(1.2)
$$J((x_t)_{t \in \mathbb{N}}, (u_t)_{t \in \mathbb{N}}) := \sum_{t=0}^{+\infty} \phi_t(x_t, u_t).$$

Now we can give the list of the considered problems of Optimal Control.

$$(\mathbf{P}_1(\sigma)): \text{ Find } ((\hat{x}_t)_{t\in\mathbb{N}}, (\hat{u}_t)_{t\in\mathbb{N}}) \in Dom(J) \cap Adm(\sigma) \text{ such that} \\ J((\hat{x}_t)_{t\in\mathbb{N}}, (\hat{u}_t)_{t\in\mathbb{N}}) \geq J((x_t)_{t\in\mathbb{N}}, (u_t)_{t\in\mathbb{N}}) \text{ for all } ((x_t)_{t\in\mathbb{N}}, (u_t)_{t\in\mathbb{N}}) \in Dom(J) \cap Adm(\sigma).$$

²⁰¹⁰ Mathematics Subject Classification. 49J21, 65K05, 39A99.

Key words and phrases. Pontryagin principle, discrete time, infinite horizon, difference equation, Banach spaces.

 $\begin{aligned} (\mathbf{P}_{2}(\sigma)): & \text{Find } ((\hat{x}_{t})_{t\in\mathbb{N}}, (\hat{u}_{t})_{t\in\mathbb{N}}) \in Adm(\sigma) \text{ such that} \\ & \lim \sup_{h\to+\infty} \sum_{t=0}^{h} (\phi_{t}(\hat{x}_{t}, \hat{u}_{t}) - \phi(x_{t}, u_{t})) \geq 0 \text{ for all } ((x_{t})_{t\in\mathbb{N}}, (u_{t})_{t\in\mathbb{N}}) \in Adm(\sigma). \\ (\mathbf{P}_{3}(\sigma)): & \text{Find } ((\hat{x}_{t})_{t\in\mathbb{N}}, (\hat{u}_{t})_{t\in\mathbb{N}}) \in Amd(\sigma) \text{ such that for all} \\ & ((x_{t})_{t\in\mathbb{N}}, (u_{t})_{t\in\mathbb{N}}) \in Adm(\sigma), \lim \inf_{h\to+\infty} \sum_{t=0}^{h} (\phi_{t}(\hat{x}_{t}, \hat{u}_{t}) - \phi(x_{t}, u_{t})) \geq 0. \end{aligned}$

These problems are classical in mathematical macroeconomic theory; cf. [10], [6], [13], [11] and references therein, and also in sustainable development theory, [8].

We study the necessary optimality conditions for these problems in the form of Pontryagin principles. The main theorem provides multipliers $(\lambda_0, (p_{t+1})_{t\in\mathbb{N}}) \in \mathbb{R}^+ \times (X^*)^{\mathbb{N}}$ not all zero such that some necessary conditions are satisfied along the optimal solution. Among the different ways to treat such a question, we choose the method of reduction to the finite horizon. This method comes

from [5] in the discrete-time framework. Notice that this viewpoint was previously used by Halkin ([7], Theorem 2.3, p. 20) in the continuous-time framework.

There exist several works on this method when X and U are finite-dimensional, cf. [6]. In the present paper we treat the case where X and U are infinite-dimensional Banach spaces. With respect to two previous papers on this question, [2] and [3], the main novelty is to avoid the use of assumptions of finiteness of the codimension of certain vector subspaces. To realize this we introduce new recursive assumptions on the partial differentials of the f_t of (1.1). We speak of recursive assumptions since they contain two successive dates t - 1 and t.

To make the reading of the paper easier, we describe the schedule of the proof of the main theorem (Theorem 2.1 below).

First step: the method of reduction to finite horizon associates to the considered problems in infinite horizon the same sequence of finite-horizon problems indexed by $h \in \mathbb{N}, h \geq 2$.

Second step: we provide conditions to ensure that we can use Multiplier Rules (in Banach spaces) on the finite-horizon problems. Hence we obtain, for each $h \in \mathbb{N}$, $h \geq 2$, a nonzero list $(\lambda_0^h, p_1^h, \dots, p_{h+1}^h) \in \mathbb{R} \times (X^*)^{h+1}$ where λ_0^h is a multiplier associated to the criterion and $(p_1^h, \dots, p_{h+1}^h)$ are multipliers associated to the (truncated) dynamical system which is transformed into a list of constraints.

Third step: we build an increasing function $\varphi : \mathbb{N} \to \mathbb{N}$ such that the subsequences $(\lambda_0^{\varphi(h)})_h$ and $(p_{t+1}^{\varphi(h)})_h$ respectively converge to λ_0 and p_{t+1} for each $t \in \mathbb{N}_*$, with $(\lambda_0, (p_{t+1})_t)$ nonzero. The Banach-Alaoglu theorem permits us to obtain weak-star convergent subsequences of $(\lambda_0^h)_h$ and $(p_{t+1}^h)_h$ for each $t \in \mathbb{N}$, and a diagonal process of Cantor permits us to obtain the same function φ for all $t \in \mathbb{N}$. The main difficulty is to avoid having $(\lambda_0, (p_{t+1})_t)$ equal to zero. Such a difficulty is due to the infinite dimension where the weak-star closure of a sphere centered at zero contains zero. To overcome this difficulty, using the Baire category theorem, we establish that a weak-star convergence implies a norm convergence on a well chosen Banach subspace of the dual space of the state space.

The main result can be adapted to the case of a dynamical system described by an inequality $x_{t+1} \leq f_t(x_t, u_t)$. To do this, X is endowed with a structure of ordered Banach space and its positive cone, denoted by X_+ is closed convex and satisfies $(X_+) \cap (-X_+) = \{0\}$. Assuming that the interior of X_+ is nonempty, it is possible to process as in [2] and [3] to obtain the conclusion of the main result (Theorem 2.1) with an additional property: $p_{t+1} \ge 0$ for all $t \in \mathbb{N}$.

Now we describe the contents of the paper. In Section 2 we present our assumptions and we give the statement of the main theorem on the Pontryagin principle. In Section 3 we recall a characterization of the closedness of the image of a linear continuous operator, a consequence of the Baire category theorem on the weak-star convergence, and we provide a diagonal process of Cantor for the weak-star convergence. In Section 4 we describe the reduction to the finite horizon and we establish consequence of our recursive assumptions on the surjectivity and on the closedness of the range of the differentials of the constraints in the finite-horizon problems. In Section 5 we give the complete proof of our main theorem.

2. The main result

First we present a list of hypotheses.

(H1): X and U are separable Banach spaces.

(H2): For all $t \in \mathbb{N}$, X_t is a nonempty open subset of X and U_t is a nonempty convex subset of U.

When $((\hat{x}_t)_{t\in\mathbb{N}}, (\hat{u}_t)_{t\in\mathbb{N}})$ is a given admissible process of one of the problems $((\mathbf{P}_i(\sigma)))$, $i \in \{1, 2, 3\}$, we consider the following conditions.

(H3): For all $t \in \mathbb{N}$, ϕ_t is Fréchet differentiable at (\hat{x}_t, \hat{u}_t) and f_t is continuously Fréchet differentiable at (\hat{x}_t, \hat{u}_t) .

(H4): For all $t \in \mathbb{N}, t \geq 2$, $D_1 f_t(\hat{x}_t, \hat{u}_t) \circ D_2 f_{t-1}(\hat{x}_{t-1}, \hat{u}_{t-1})(U) + D_2 f_t(\hat{x}_t, \hat{u}_t)(T_{U_t}(\hat{u}_t)) = X.$ (H5): $D_1 f_1(\hat{x}_1, \hat{u}_1) \circ D_2 f_0(\hat{x}_0, \hat{u}_0)(T_{U_0}(\hat{u}_0)) + D_2 f_1(\hat{x}_1, \hat{u}_1)(T_{U_1}(\hat{u}_1)) = X.$ (H6): $ri(T_{U_0}(\hat{u}_0)) \neq \emptyset$ and $ri(T_{U_1}(\hat{u}_1)) \neq \emptyset.$

In (H3), since U_t is not necessarily a neighborhood of \hat{u}_t , the meaning of this condition is that there exist an open neighborhood V_t of (\hat{x}_t, \hat{u}_t) in $X \times U$ and a Fréchet differentiable function (respectively continuously Fréchet differentiable mapping) $\tilde{\phi}_t : V_t \to \mathbb{R}$ (respectively $\tilde{f}_t : V_t \to X$) such that $\tilde{\phi}_t$ and ϕ_t (respectively \tilde{f}_t and f_t) coincide on $V_t \cap (X_t \times U_t)$. Moreover D_1 and D_2 denotes the partial Fréchet differentials with respect to the first (vector) variable and with respect to the second (vector) variable respectively. About (H4), (H5) and (H6), when A is a convex subset of U, $\hat{u} \in A$, the set $T_A(\hat{u})$ is the closure of $\mathbb{R}_+(A - \hat{u})$; it is called the tangent cone of A at \hat{u} as it is usually defined in Convex Analysis, [1] p. 166. About (H6), if $\operatorname{aff}(T_{U_t}(\hat{u}_t))$ denotes the affine hull of $T_{U_t}(\hat{u}_t)$, $ri(T_{U_t}(\hat{u}_t))$ denotes the (relative) interior of $T_{U_t}(\hat{u}_t)$ in $\operatorname{aff}(T_{U_t}(\hat{u}_t))$. Such definition of the relative interior of a convex is given in [12], p. 14-15, where it is denoted by *rint*.

Now we state the main result of the paper.

Theorem 2.1. Let $((\hat{x}_t)_{t\in\mathbb{N}}, (\hat{u}_t)_{t\in\mathbb{N}})$ be an optimal process for one of the problems $(\mathbf{P}_i(\sigma)), i \in \{1, 2, 3\}$. Under (H1-H6), there exist $\lambda_0 \in \mathbb{R}$ and $(p_{t+1})_{t\in\mathbb{N}} \in (X^*)^{\mathbb{N}}$ which satisfy the following conditions.

- (1) $(\lambda_0, p_1, p_2) \neq (0, 0, 0).$
- (2) $\lambda_0 \ge 0$.
- (3) $p_t = p_{t+1} \circ D_1 f_t(\hat{x}_t, \hat{u}_t) + \lambda_0 D_1 \phi_t(\hat{x}_t, \hat{u}_t), \text{ for all } t \in \mathbb{N}, t \ge 1.$
- (4) $\langle \lambda_0 D_2 \phi_t(\hat{x}_t, \hat{u}_t) + p_{t+1} \circ D_2 f_t(\hat{x}_t, \hat{u}_t), u_t \hat{u}_t \rangle \leq 0$, for all $u_t \in U_t$, for all $t \in \mathbb{N}$.

In comparison with Theorem 2.2 in [3], note that

- assumptions (H1), (H2), (H3) are essentially identical to assumptions (A1), (A2), (A3) of [3],
- assumptions (A4) and (A5) of [3] are replaced by (H4) and (H5) to permits us to delete a condition of finiteness of a codimension,
- assumption (A6) of [3] is replaced by (H6) which also permits to avoid a condition of finiteness of a codimension.

Notice that (H6) is automatically satisfied when the dimension of U is finite.

3. Functional analytic results

In this section, first we recall a characterization of the closedness of the image of a linear continuous operator. Secondly we state a result which is a consequence of the Baire category theorem. After this, we give a version of the diagonal process of Cantor for the weak-star convergence.

Proposition 3.1. Let E and F be Banach spaces, and $L \in \mathfrak{L}(E, F)$ (the space of all linear continuous mappings from E into F). The two following assertions are equivalent.

- (i) ImL is closed in F.
- (ii) There exists $c \in (0, +\infty)$ such that for all $y \in ImL$, there exists $x_y \in E$ verifying $Lx_y = y$ and $||y|| \ge c||x_y||$.

This result is proven in [2] (Lemma 3.4) and in [4] (Lemma 2.1).

Proposition 3.2. Let Y be a real Banach space; Y^* is its topological dual space. Let $(\pi_h)_{h\in\mathbb{N}} \in (Y^*)^{\mathbb{N}}$ and $(\rho_h)_{h\in\mathbb{N}} \in (\mathbb{R}_+)^{\mathbb{N}}$. Let K be a nonempty closed convex subset of Y such that $ri(K) \neq \emptyset$. Let $a \in K$ and we set $S := \overline{\operatorname{aff}}(K) - a$ which is a Banach subspace. We assume that the following conditions are fulfilled.

- (1) $\rho_h \to 0$ when $h \to +\infty$.
- (2) $\pi_h \xrightarrow{w^*} 0$ (weak-star convergence) when $h \to +\infty$.

(3) For all $y \in K$, there exists $c_y \in \mathbb{R}$ such that $\pi_h(y) \leq c_y \rho_h$ for all $h \in \mathbb{N}$.

Then we have $\|\pi_{h|_S}\|_{S^*} \to 0$ when $h \to +\infty$.

This result is established in [3] (Proposition 3.5) where several consequences and generalizations are provided. In the following result, when $t \in \mathbb{N}$, we set $[t, +\infty)_{\mathbb{N}} := [t, +\infty) \cap \mathbb{N}$ and $\mathbb{N}_* := [1, +\infty)_{\mathbb{N}}$. **Proposition 3.3.** Let Y be a real Banach space; Y^* is its topological dual space. For every $(t,h) \in \mathbb{N} \times \mathbb{N}_*$ such that $t \leq h$ we consider an element $\pi_{t+1}^h \in Y^*$. We assume that, for every $t \in \mathbb{N}$, the sequence $(\pi_{t+1}^h)_{h \in [t,+\infty)_{\mathbb{N}}}$ is bounded in Y^* . Then there exists an increasing function $\beta : \mathbb{N}_* \to \mathbb{N}_*$ such that, for all $t \in \mathbb{N}$, there exists $\overline{\pi}_{t+1} \in Y^*$ verifying $\pi_{t+1}^{\beta(h)} \xrightarrow{w^*} \overline{\pi}_{t+1}$ when $h \to +\infty$.

Proof. Using the Banach-Alaoglu theorem, since $(\pi_1^h)_{h\in[0,+\infty)_{\mathbb{N}}}$ is bounded in Y^* , there exists an increasing function $\alpha_1 : [0,+\infty)_{\mathbb{N}} \to [0,+\infty)_{\mathbb{N}}$ and $\overline{\pi}_1 \in Y^*$ such that $\pi_1^{\alpha_1(h)} \stackrel{w^*}{\to} \overline{\pi}_1$ when $h \to +\infty$. Using the same argument, since $(\pi_2^{\alpha_1(h)})_{h\in[1,+\infty)_{\mathbb{N}}}$ is bounded, there exists an increasing function $\alpha_2 : [1,+\infty)_{\mathbb{N}} \to [1,+\infty)_{\mathbb{N}}$ and $\overline{\pi}_2 \in Y^*$ such that $\pi_2^{\alpha_1 \circ \alpha_2(h)} \stackrel{w^*}{\to} \overline{\pi}_2$ when $h \to +\infty$. Iterating the reasoning, for every $t \in \mathbb{N}_*$, there exist an increasing function $\alpha_t : [t,+\infty)_{\mathbb{N}} \to [t,+\infty)_{\mathbb{N}}$ and $\overline{\pi}_{t+1} \in Y^*$ such that $\pi_{t+1}^{\alpha_1 \circ \ldots \circ \alpha_t(h)} \stackrel{w^*}{\to} \overline{\pi}_{t+1}$ when $h \to +\infty$. We define the function $\beta : [0,+\infty)_{\mathbb{N}} \to [0,+\infty)_{\mathbb{N}}$ by setting $\beta(h) := \alpha_1 \circ \ldots \circ \alpha_h(h)$. we arbitrarily fix $t \in \mathbb{N}_*$ and we define the function $\delta_t : [t,+\infty)_{\mathbb{N}} \to [t,+\infty)_{\mathbb{N}}$ by setting $\delta_t(t) := t$ and $\delta_t(h) := \alpha_{t+1} \circ \ldots \circ \alpha_h(h)$ when h > t. When h = t, we have $\delta_t(t+1) = \alpha_{t+1}(t+1) \ge t+1 > t = \delta_t(t)$. When $h \in [t+1,+\infty)_{\mathbb{N}}$, we have $\alpha_{t+1}(h+1) \ge h+1 > h$ which implies

$$\delta_t(h+1) = (\alpha_{t+1} \circ \dots \circ \alpha_h)(\alpha_{t+1}(h+1)) > (\alpha_{t+1} \circ \dots \circ \alpha_h)(h) = \delta_t(h)$$

since $(\alpha_{t+1} \circ ... \circ \alpha_h)$ is increasing. Hence we have proven that δ_t is increasing. Since $\beta_{|_{[t,+\infty)_{\mathbb{N}}}} = (\alpha_1 \circ ... \circ \alpha_t) \circ \delta_t$, we can say that $(\pi_{t+1}^{\beta(h)})_{h \in [t,+\infty)_{\mathbb{N}}}$ is a subsequence of $(\pi_{t+1}^{\alpha_1 \circ ... \circ \alpha_t(h)})_{h \in [t,+\infty)_{\mathbb{N}}}$, we obtain $\pi_{t+1}^{\beta(h)} \xrightarrow{w^*} \overline{\pi}_{t+1}$ when $h \to +\infty$.

4. REDUCTION TO THE FINITE HORIZON

Let $((\hat{x}_t)_{t\in\mathbb{N}}, (\hat{u}_t)_{t\in\mathbb{N}})$ be an optimal process for one of the problems $(\mathbf{P}_i(\sigma)), i \in \{1, 2, 3\}$. The method of reduction to finite horizon consists of considering a sequence of finite-horizon problems as follows:

$$(\mathbf{F}_{h}(\sigma)) \begin{cases} \text{Maximize} & J_{h}(x_{1}, \dots, x_{h}, u_{0}, \dots, u_{h}) := \sum_{t=0}^{h} \phi_{t}(x_{t}, u_{t}) \\ \text{when} & (x_{t})_{1 \le t \le h} \in \prod_{t=1}^{h} X_{t}, (u_{t})_{0 \le t \le h} \in \prod_{t=0}^{h} U_{t} \\ & \forall t \in \{0, \dots, h\}, x_{t+1} = f_{t}(x_{t}, u_{t}) \\ & x_{0} = \sigma, x_{h+1} = \hat{x}_{t+1} \end{cases}$$

The proof of the following lemma is given in [5].

Lemma 4.1. When $((\hat{x}_t)_{t\in\mathbb{N}}, (\hat{u}_t)_{t\in\mathbb{N}})$ is an optimal process for one of the problems $(\mathbf{P}_i(\sigma)), i \in \{1, 2, 3\},$ then, for all $h \in \mathbb{N}_*, (\hat{x}_1, ..., \hat{x}_h, \hat{u}_0, ..., \hat{u}_h)$ is an optimal solution of $(\mathbf{F}_h(\sigma))$.

Notice that this result does not need any special assumption. Now we introduce notation to work on these problems. We write $\mathbf{x}^h := (x_1, ..., x_h) \in \prod_{t=1}^h X_t$, $\mathbf{u}^h := (u_0, ..., u_h) \in \prod_{t=0}^h U_t$. For all $h \in \mathbb{N}_*$ and for all $t \in \mathbb{N}$, we introduce the mapping

$$g_t^h : (\prod_{t=1}^h X_t) \times (\prod_{t=0}^h U_t) \to X_{t+1} \text{ by setting}$$

$$(4.1) \qquad g_t^h(\mathbf{x}^h, \mathbf{u}^h) := \begin{cases} -x_1 + f_0(\sigma, u_0) & \text{if } t = 0\\ -x_{t+1} + f_t(x_t, u_t) & \text{if } t \in \{1, ..., h-1\}\\ -\hat{x}_{h+1} + f_h(x_h, u_h). \end{cases}$$

We introduce the mapping $g^h: (\prod_{t=1}^h X_t) \times (\prod_{t=0}^h U_t) \to X^{h+1}$ defined by $g^{h}(\mathbf{x}^{h}, \mathbf{u}^{h}) := (g^{h}_{0}(\mathbf{x}^{h}, \mathbf{u}^{h}), \dots, g^{h}_{h}(\mathbf{x}^{h}, \mathbf{u}^{h})).$ (4.2)

Under (H3), g^h is of class C^1 . We introduce the following conditions on the differentials of the f_t .

(4.3)
$$\forall t \in \mathbb{N}, \ ImDf_t(\hat{x}_t, \hat{u}_t) \text{ is closed in } X.$$

$$\forall t \in \mathbb{N}_*, Im(D_1 f_t(\hat{x}_t, \hat{u}_t) \circ D_2 f_{t-1}(\hat{x}_{t-1}, \hat{u}_{t-1})) + ImD_2 f_t(\hat{x}_t, \hat{u}_t) = ImDf_t(\hat{x}_t, \hat{u}_t).$$

(4.5)
$$\forall t \in \mathbb{N}, t \ge 2, \quad ImDf_t(\hat{x}_t, \hat{u}_t) = X.$$

(4.6)
$$Im(D_1f_1(\hat{x}_1, \hat{u}_1) \circ D_2f_0(\sigma, \hat{u}_0)) + ImD_2f_1(\hat{x}_1, \hat{u}_1) = X.$$

Lemma 4.2. We assume that (H3) is fulfilled.

- (i) Under (4.3) and (4.4), $ImDg^h(\mathbf{x}^h, \mathbf{u}^h)$ is closed.
- (ii) Under (4.5) and (4.6), $Dg^h(\mathbf{x}^h, \mathbf{u}^h)$ is surjective.

Proof. (i) To abridge the writing we set $D\hat{f}_t := Df_t(\hat{x}_t, \hat{u}_t)$ and $D_i\hat{f}_t := D_if_t(\hat{x}_t, \hat{u}_t)$ when $i \in \{1, 2\}$. The condition (H3) implies that g^h is Fréchet differentiable at $(\mathbf{x}^h, \mathbf{u}^h).$

We arbitrarily fix $\mathbf{z}^h = (z_0, ..., z_h) \in ImDg^h(\mathbf{x}^h, \mathbf{u}^h)$. Therefore there exist $\mathbf{y}^{h,0} = (y_1^0, ..., y_h^0) \in X^h$ and $\mathbf{v}^{h,0} = (v_0^0, ..., v_h^0) \in U^{h+1}$ such that $\mathbf{z}^h = Dg^T(\mathbf{x}^h, \mathbf{u}^h)(\mathbf{y}^{h,0}, \mathbf{v}^{h,0})$ which is equivalent to the set of the three following

equations

(4.7)
$$-y_1^0 + D_2 f_0(\sigma, \hat{u}_0) v_0^0 = z_0$$

(4.8)
$$\forall t \in \{1, ..., h-1\}, \quad -y_{t+1}^0 + D_1 \hat{f}_t y_t^0 + D_2 \hat{f}_t v_t^0 = z_t$$

(4.9)
$$D_1 \hat{f}_h y_h^0 + D_2 \hat{f}_h v_h^0 = z_h$$

We introduce the linear continuous operator $L_0 \in \mathfrak{L}(X \times U, X)$ by setting

(4.10)
$$L_0(y_1, v_0) := -y_1 + D_2 \tilde{f}_0 v_0.$$

Notice that L_0 is surjective since $L_0(X \times \{0\}) = X$; therefore ImL_0 is closed in X. From (4.7) we have $z_0 \in ImL_0$. Using Proposition 3.1 on L_0 we know that

$$\begin{cases} \exists a_0 \in (0, +\infty), \forall z_0 \in X, \exists y_1^* \in X, \exists v_0^* \in U \text{ s.t. } L_0(y_1^*, v_0^*) = z_0 \\ \text{and} \max\{\|y_1^*\|, \|v_0^*\|\} \le a_0 \cdot \|z_0\| \end{cases}$$

i.e. we have proven

(4.11)
$$\exists a_0 \in (0, +\infty), \exists y_1^* \in X, \exists v_0^* \in U \text{ s.t.} \\ -y_1^* + D_2 \hat{f}_0 v_0^* = z_0 \text{ and } \max\{\|y_1^*\|, \|v_0^*\|\} \le a_0 \cdot \|z_0\| \}$$

It is important to notice that a_0 does not depend on z_0 .

We introduce the linear continuous operator $L_1 \in \mathfrak{L}(X \times U, X)$ by setting

$$L_1(y_2, v_1) := -y_2 + D_2 f_1 v_1$$

Since $L_1(X \times \{0\}) = X$, L_1 is surjective and hence $z_1 - D_1 \hat{f}_1 y_1^* \in ImL_1$. Using Proposition 3.1 on L_1 , we obtain

$$\begin{cases} \exists b_1 \in (0, +\infty), \exists y_2^* \in X, \exists v_1^* \in U \text{ s.t.} \\ L_1(y_2^*, v_1^*) = z_1 - D_1 \hat{f}_1 y_1^* \text{ and} \\ \max\{\|y_2^*\|, \|v_1^*\|\} \le b_1 \cdot \|z_1 - D_1 \hat{f}_1 y_1^*\|. \end{cases}$$

Using (4.11) we deduce from the last inequality

$$\max\{\|y_2^*\|, \|v_1^*\|\} \leq b_1 \cdot (\|z_1\| + \|D_1\hat{f}_1\| \cdot \|y_1^*\|) \\ \leq b_1 \cdot (\|z_1\| + \|D_1\hat{f}_1\| \cdot a_0 \cdot \|z_0\|) \\ \leq b_1 \cdot (1 + a_0 \cdot \|D_1\hat{f}_1\|) \cdot \max\{\|z_0\|, \|z_1\|\}.$$

We set $a_1 := \max\{a_0, b_1 \cdot (1 + a_0 \cdot ||D_1\hat{f}_1||)\}$, and then we have proven the following assertion.

(4.12)
$$\begin{aligned} \exists a_1 \in (0, +\infty), \exists (y_1^*, y_2^*, v_0^*, v_1^*) \in X^2 \times U^2 \text{ s.t.} \\ -y_1^* + D_2 \hat{f}_0 v_0^* = z_0, \quad -y_2^* + D_1 \hat{f}_1 y_1^* + D_2 \hat{f}_1 v_1^* = z_1, \\ \max\{|y_1^*\|, \|y_2^*\|, \|v_0^*\|, \|v_1^*\|\} \le a_1 \cdot \max\{\|z_0\|, \|z_1\|\}. \end{aligned}$$

It is important to notice that a_1 does not depend on z_0 , z_1 . We iterate the reasoning until h-2 and we obtain

$$\left. \begin{array}{l} \exists a_{h-2} \in (0, +\infty), \exists (y_t^*)_{1 \le t \le h-1} \in X^{h-1}, \exists (v_t^*)_{0 \le t \le h-2} \in U^{h-1} \text{ s.t.} \\ (4.13) \quad -y_1^* + D_2 \hat{f}_0 v_0^* = z_0, \quad \forall t \in \{1, ..., h-2\}, -t_{t+1}^* + D_1 \hat{f}_t y_t^* + D_2 \hat{f}_t v_t^* = z_t \\ \max\{\max_{1 \le t \le h-1} \|y_t^*\|, \max_{0 \le t \le h-2} \|v_t^*\|\} \le a_{h-2} \max_{0 \le t \le h-2} \|z_t\|. \end{array} \right\}$$

From (4.9) we know that $z_h \in ImD\hat{f}_h$. Moreover we have

$$D_1\hat{f}_h z_{h-1} \subset ImD\hat{f}_h$$
 and $D_1\hat{f}_h \circ D_1\hat{f}_{h-1}y_{h-1}^* \in ImD_1\hat{f}_h \subset ImD\hat{f}_h$

and therefore we have

(4.14)
$$z_h + D_1 \hat{f}_{h-1} z_{h-1} - D_1 \hat{f}_h \circ D_1 \hat{f}_{h-1} y_{h-1}^* \in ImD\hat{f}_h$$

Introduce the linear continuous operator $\Lambda \in \mathfrak{L}(U \times U, X)$ by setting

(4.15)
$$\Lambda(v,w) := D_1 \hat{f}_h \circ D_2 \hat{f}_{h-1} v + D_2 \hat{f}_h w$$

Under assumptions (4.4) and (4.5) we have $Im\Lambda = ImD\hat{f}_h$ and $Im\Lambda$ is closed in X. After (4.14), using Proposition 3.1 on Λ we obtain

$$(4.16) \quad \begin{aligned} \exists c \in (0, +\infty), \exists (v_{h-1}^*, v_h^*) \in U \times U, \text{ s.t.} \\ & \Lambda(v_{h-1}^*, v_h^*) = z_h + D_1 \hat{f}_h z_{h-1} - D_1 \hat{f}_h \circ D_1 \hat{f}_{h-1} y_{h-1}^* \\ & \text{i.e.} \\ & D_1 \hat{f}_h \circ D_2 \hat{f}_{h-1} v_{h-1}^* + D_2 \hat{f}_h v_h^* = \\ & z_h + D_1 \hat{f}_h z_{h-1} - D_1 \hat{f}_h \circ D_1 \hat{f}_{h-1} y_{h-1}^* \text{ and} \\ & \max\{ \|v_{h-1}^*\|, \|v_h^*\|\} \le c \cdot \|z_h + D_1 \hat{f}_h z_{h-1} - D_1 \hat{f}_h \circ D_1 \hat{f}_{h-1} y_{h-1}^* \|. \end{aligned} \right\}$$

M. BACHIR AND J. BLOT

From this last inequality, using (4.13), we obtain

 $\begin{aligned} \max\{\|v_{h-1}^*\|, \|v_h^*\|\} \\ &\leq c \cdot (\|z_h\| + \|D_1\hat{f}_h\| \cdot \|z_{h-1}\| + \|D_1\hat{f}_h \circ D_1\hat{f}_{h-1}\| \cdot \|y_{h-1}^*\|) \\ &\leq c \cdot (\|z_h\| + \|D_1\hat{f}_h\| \cdot \|z_{h-1}\| + \|D_1\hat{f}_h \circ D_1\hat{f}_{Th1}\| \cdot a_{h-2} \cdot \max_{1 \leq t \leq h-2} \|z_t\|) \\ &\leq c \cdot (1 + \|D_1\hat{f}_h\| + a_{h-2} \cdot \|D_1\hat{f}_h \circ D_1\hat{f}_{h-1}\|) \cdot \max_{1 \leq t \leq h} \|z_t\|. \end{aligned}$

We set $c_1 := c \cdot (1 + ||D_1\hat{f}_h|| + a_{h-2} \cdot ||D_1\hat{f}_h \circ D_1\hat{f}_{h-1}||) \in (0, +\infty)$. Then we have proven the following assertion.

(4.17)
$$\exists c_1 \in (0, +\infty), \max\{\|v_{h-1}^*\|, \|v_h^*\|\} \le c_1 \cdot \max_{1 \le t \le h} \|z_t\|.$$

We set

(4.18)
$$y_h^* := D_2 \hat{f}_{h-1} v_{h-1}^* + D_1 \hat{f}_{h-1} y_{h-1}^* - z_{h-1}.$$

This equality implies

(4.19)
$$-y_h^* + D_1 \hat{f}_{h-1} y_{h-1}^* + D_2 \hat{f}_{h-1} v_{h-1}^* = z_{h-1}$$

which is the penultimate wanted equation.

Notice that we have $||y_h^*|| \le ||D_2 \hat{f}_{h-1}|| \cdot ||v_{h-1}^*|| + ||D_1 \hat{f}_{h-1}|| \cdot ||y_{h-1}^*|| + ||z_{h-1}||$, and using (4.17) and (4.18) we obtain

$$\begin{aligned} \|y_h^*\| &\leq \|D_2 f_{h-1}\| \cdot c_1 \cdot \max_{1 \leq t \leq h} \|z_t\| \\ &+ \|D_1 \hat{f}_{h-1}\| \cdot a_{h-2} \cdot \max_{1 \leq t \leq h-2} \|z_t\| + \|z_{h-1}\| \\ &\leq (c_1 \cdot \|D_2 \hat{f}_{h-1}\| + a_{h-2} \cdot \|D_1 \hat{f}_{h-1}\| + 1) \cdot \max_{1 \leq t \leq h} \|z_t\|. \end{aligned}$$

We set $c_2 := c_1 \cdot \|D_2 \hat{f}_{h-1}\| + a_{h-2} \cdot \|D_1 \hat{f}_{h-1}\| + 1$, and so we have proven

(4.20)
$$\exists c_2 \in (0, +\infty), \|y_h^*\| \le c_2 \cdot \max_{1 \le t \le h} \|z_t\|.$$

We set $a_h := \max\{a_{h-3}, c_1, c_2\}$, and from (4.13), (4.17) and (4.20) we have proven

(4.21)
$$\exists a_h \in (0, +\infty), \max\{\max_{1 \le t \le h} \|y_t^*\|, \max_{0 \le t \le h} \|v_t^*\|\} \le a_h \cdot \max_{1 \le t \le h} \|z_t\|$$

Now we show that the last equation is satisfied by y_h^* and v_h^* . Using (4.18) and (4.16), we obtain

$$\begin{split} &D_1 \hat{f}_h y_h^* + D_2 \hat{f}_h v_h^* \\ &= D_1 \hat{f}_T (D_2 \hat{f}_{h-1} v_{h-1}^* + D_1 \hat{f}_{h-1} y_{h-1}^* - z_{h-1}) + D_2 \hat{f}_h v_h^* \\ &= (D_1 \hat{f}_h \circ (D_2 \hat{f}_{h-1} v_{h-1}^* + D_2 \hat{f}_h v_h^*) + D_1 \hat{f}_h \circ D_1 \hat{f}_{h-1} y_{h-1}^* - D_1 \hat{f}_h z_{h-1} \\ &= (z_h + D_1 \hat{f}_h z_{h-1} - D_1 \hat{f}_h \circ D_1 \hat{f}_{h-1} y_{h-1}^*) + D_1 \hat{f}_h \circ D_1 \hat{f}_{h-1} y_{h-1}^* - D_1 \hat{f}_h z_{h-1} \\ &= z_h. \end{split}$$

We have proven that

(4.22)
$$D_1 \hat{f}_h y_h^* + D_2 \hat{f}_h v_h^* = z_h$$

28

From (4.13), (4.19), (4.21) and (4.22) we have proven the following assertion

$$\begin{cases} \exists a_h \in (0, +\infty), \forall (z_t)_{0 \le t \le h} \in ImDg_h(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h), \\ \exists (y_t^*)_{1 \le t \le h} \in X^h, \exists (v_t^*)_{0 \le t \le h} \in U^{h+1}, \text{ such that} \\ -y_1^* + D_2 \hat{f}_0 v_0^* = z_0, \forall t \in \{1, \dots, h-1\}, \quad -y_{t+1} + D_1 \hat{f}_t y_t^* + D_2 \hat{f}_t v_t^* = z_t, \\ D_1 \hat{f}_h y_h^* + D_2 \hat{f}_h v_h^* = z_h, \text{ and} \\ \max\{\max_{1 \le t \le h} \|y_t^*\|, \max_{0 \le t \le h} \|v_t^*\|\} \le a_h \cdot \max_{1 \le t \le h} \|z_t\|. \end{cases}$$

This last assertion is equivalent to the following one

$$\begin{cases} \exists a_h \in (0, +\infty), \forall \mathbf{z}^h = (z_t)_{0 \le t \le h} \in ImDg^h(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h), \\ \exists \mathbf{y}^{h,*} = (y_t^*)_{1 \le t \le h} \in X^h, \exists \mathbf{v}^{h,*} = (v_t^*)_{0 \le t \le h} \in U^{h+1}, \text{ such that} \\ Dg^h(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h)(\mathbf{y}^{h,*}, \mathbf{v}^{h,*}) = \mathbf{z}^h \text{ and } \|(\mathbf{y}^{h,*}, \mathbf{v}^{h,*})\| \le a_h \cdot \|\mathbf{z}^h\|. \end{cases}$$

Now using Proposition 3.1 on the operator $Dg^h(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h)$, the previous assertion permits us to assert that $ImDg^h(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h)$ is closed in X^{h+1} , and the proof of (i) is complete.

(ii) We arbitrarily fix $\mathbf{z}^h = (z_1, ..., z_h) \in X^h$. Since $D\hat{f}_h$ is surjective, there exists $y_h^{\#} \in X$ and $v_h^{\#} \in U$ such that $D\hat{f}_h(y_h^{\#}, v_h^{\#}) = z_h$. Since $D\hat{f}_{h-1}$ is surjective, there exists $y_{h-1}^{\#} \in X$ and $v_{h-1}^{\#} \in U$ such that $D\hat{f}_{h-1}(y_{h-1}^{\#}, v_{h-1}^{\#}) = z_{h-1} + y_h^{\#}$. We iterate this backward reasoning until t = 2 to obtain

(4.23)
$$\begin{cases} \forall t \in \{2, ..., h\}, \exists (y_t^{\#}, v_t^{\#}) \in X \times U \text{ s.t. } D\hat{f}_h(y_h^{\#}, v_h^{\#}) = z_h \\ \text{and } \forall t \in 2, ..., h - 1, -y_t^{\#} + D\hat{f}_t(y_t^{\#}, v_t^{\#}) = z_t. \end{cases}$$

Now we introduce the linear continuous operator $M \in \mathfrak{L}(U \times U, X)$ by setting $M(v_0, v_1) := D_1 \hat{f}_1 \circ D_2 \hat{f}_0 v_0 + D_2 \hat{f}_1 v_1$. From (4.6) we have ImM = X i.e. M is surjective. Therefore we obtain

(4.24)
$$\exists (v_0^{\#}, v_1^{\#}) \in U \times U \text{ s.t. } D_1 \hat{f}_1 \circ D_2 \hat{f}_0 v_0^{\#} + D_2 \hat{f}_1 v_1^{\#} = z_1 + y_2^{\#} + D_1 \hat{f}_1 z_0.$$

We set $y_1^{\#} := D_2 \hat{f}_0 v_0^{\#} - z_0$. Hence we obtain

(4.25)
$$-y_1^{\#} + D_2 \hat{f}_0 v_0^{\#} = z_0.$$

Using (4.24) and (4.25), we calculate

$$\begin{aligned} -y_2^{\#} + D_1 \hat{f}_1 y_1^{\#} + D_2 \hat{f}_1 v_1^{\#} &= -y_2^{\#} + D_1 \hat{f}_1 (D_2 \hat{f}_0 v_0^{\#} - z_0) + D_2 \hat{f}_1 v_1^{\#} \\ &= -y_2^{\#} + (D_1 \hat{f}_1 \circ D_2 \hat{f}_0 v_0^{\#} + D_2 \hat{f}_1 v_1^{\#}) - D_1 \hat{f}_1 z_0 \\ &= -y_2^{\#} + (z_1 + y_2^{\#} + D_1 \hat{f}_1 z_0) - D_1 \hat{f}_1 z_0 = z_1. \end{aligned}$$

We have proven

(4.26)
$$-y_2^{\#} + D_1 \hat{f}_1 y_1^{\#} + D_2 \hat{f}_1 v_1^{\#} = z_1 \cdot z_1 \cdot z_1 \cdot z_1 \cdot z_2 \cdot z_1 \cdot z_1 \cdot z_1 \cdot z_2 \cdot z_1 \cdot z_1$$

From (4.23), (4.25) and (4.26) we have proven

$$(4.27) \qquad \begin{array}{l} \forall (z_t)_{0 \le t \le h} \in X^{h+1}, \exists (y_t^{\#})_{1 \le t \le h} \in X^h, \exists (v_t^{\#})_{0 \le t \le h} \in U^{h+1} \text{ s.t.} \\ -y_1^{\#} + D_2 \hat{f}_0 v_0^{\#} = z_0, \forall t \in \{1, ..., h-1\} \quad -y_{t+1}^{\#} + D\hat{f}(y_t^{\#}, v_t^{\#}) = z_t \\ \text{and} \quad D\hat{f}_h(y_h^{\#}, v_h^{\#}) = z_h. \end{array} \right\}$$

This assertion is equivalent to

$$\forall \mathbf{z}^h \in X^{h+1}, \exists \mathbf{y}^{h,\#} \in X^h, \exists \mathbf{v}^{h,\#} \in U^{h+1} \text{ s.t. } Dg^h(\mathbf{x}^h, \mathbf{u}^h)(\mathbf{y}^{h,\#}, \mathbf{v}^{h,\#}) = \mathbf{z}^h$$

which means that $Dg^h(\mathbf{x}^h, \mathbf{u}^h)$ is surjective.

Lemma 4.3. Let $((\hat{x}_t)_{t\in\mathbb{N}}, (\hat{u}_t)_{t\in\mathbb{N}})$ be an optimal solution of one of the problems $(\mathbf{P}_{i}(\sigma)), i \in \{1, 2, 3\}.$ Under (H1), (H2), (H3), (4.3) and (4.4), for all $h \in \mathbb{N}_{*}$, there exist $\lambda_0^h \in \mathbb{R}$ and $(p_{t+1}^h)_{0 \le t \le h} \in (X^*)^{h+1}$ such that the following assertions hold.

- (a) λ_0^h and $(p_{t+1}^h)_{0 \le t \le h}$ are not simultaneously equal to zero.

- (b) $\lambda_0^h \ge 0.$ (c) $p_t^h = p_{t+1}^h \circ D_1 f_t(\hat{x}_t, \hat{u}_t) + \lambda_0^h D_1 \phi_t(\hat{x}_t, \hat{u}_t) \text{ for all } t \in \mathbb{N}_*.$ (d) $\langle \lambda_0^h D_2 \phi_t(\hat{x}_t, \hat{u}_t) + p_{t+1}^h \circ D_2 f_t(\hat{x}_t, \hat{u}_t), u_t \hat{u}_t \rangle \le 0 \text{ for all } t \in \{0, ..., h\}, \text{ for } t \in \{0, ..., h\},$ all $u_t \in U_t$.

Moreover, for all $h \ge 2$, if in addition we assume (H4), (H5) and (H6) fulfilled, the following assertions hold.

- (e) For all $t \in \{1, ..., h+1\}$, there exists $a_t, b_t \in \mathbb{R}_+$ such that, for all $s \in \mathbb{R}_+$ $\{1, ..., h\}, \|p_t^h\| \le a_t \lambda_0^h + b_t \|p_s^h\|.$
- (f) For all $t \in \{1, ..., h\}$, $(\lambda_0^h, p_t^h) \neq (0, 0)$.
- (g) For all $t \in \{1, ..., h\}$, for all $z \in A_t := D_2 f_{t-1}(\hat{x}_{t-1}, \hat{u}_{t-1})(T_{U_{t-1}}(\hat{u}_{t-1})),$ there exists $c_z \in \mathbb{R}$ such that $p_t^h(z) \leq c_z \lambda_0^h$ for all $h \geq t$.

Proof. Let $h \in \mathbb{N}_*$. Using Lemma 4.1, (4.1) and (4.2), we know that $(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h)$ (where $\hat{\mathbf{x}}^h = (x_1^h, ..., x_h^h)$ and $\hat{\mathbf{u}}^h = (u_0^h, ..., u_h^h)$, is an optimal solution of the following maximization problem,

$$\begin{array}{ll} \text{Maximize} & J_h(\mathbf{x}^h, \mathbf{u}^h) \\ \text{when} & (\mathbf{x}^h, \mathbf{u}^h) \in (\prod_{t=1}^h X_t) \times (\prod_{t=0}^h U_t), \\ & g^h(\mathbf{x}^h, \mathbf{u}^h) = 0. \end{array}$$

From (H3) we know that J_h is Fréchet differentiable at $(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h)$ and g^h is Fréchet continuously differentiable at $(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h)$. From (4.3), (4.4) and Lemma 4.2 we know that $ImDq^h(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h)$ is closed in X^{h+1} . Now using the multiplier rule which is given in [9] (Theorem 3.5 p. 106–111 and Theorem 5.6 p. 118) and explicitly written in [3] (Theorem 4.4), and proceeding as in the proof of Lemma 4.5 of [3], we obtain the assertions (a), (b), (c), (d).

The proof of assertions (e), (f), (g) is given by Lemma 4.7 of [3]. The proof of this Lemma 4.7 uses the condition $0 \in Int[Df(\hat{x}_t, \hat{u}_t)(X \times T_{U_t}(\hat{u}_t)) \cap B_{X \times U}]$ where $B_{X \times U}$ is the closed unit ball of $X \times U$. It suffices to notice that our assumption (H4) implies this condition.

Remark 4.4. In Lemma 4.5 of [3] the finiteness of the codimension of $ImD_2f(\hat{x}_t, \hat{u}_t)$ is useful to ensure the closedness of $ImDq^h(\hat{\mathbf{x}}^h, \hat{\mathbf{u}}^h)$. Here we can avoid this assumption of finiteness thanks to the recursive assumptions.

The following proposition is used in the proof of the main result.

Proposition 4.5. Let $(\hat{x}_t)_{t \in \mathbb{N}}, (\hat{u}_t)_{t \in \mathbb{N}})$ be an optimal solution of one of the problems $(\mathbf{P}_i(\sigma)), i \in \{1, 2, 3\}$. Under (H1-H6) we introduce

$$Z_0 := D_2 f_0(\sigma, \hat{u}_0)(T_{U_0}(\hat{u}_0))$$
 and $Z_1 := D_2 f_1(\hat{x}_1, \hat{u}_1)(T_{U_1}(\hat{u}_1)).$

Then, for all $h \in \mathbb{N}$, $h \ge 2$, there exist $\lambda_0^h \in \mathbb{R}$ and $(p_{t+1}^h)_{0 \le t \le h} \in (X^*)^{h+1}$ such that the following assertions hold.

- $\begin{array}{ll} (1) \ \lambda_0^h \geq 0. \\ (2) \ p_t^h = p_{t+1}^h \circ D_1 f_t(\hat{x}_t, \hat{u}_t) + \lambda_0^h D_1 \phi_t(\hat{x}_t, \hat{u}_t) \ for \ all \ t \in \mathbb{N}_*. \\ (3) \ \langle \lambda_0^h D_2 \phi_t(\hat{x}_t, \hat{u}_t) + p_{t+1}^h \circ D_2 f_t(\hat{x}_t, \hat{u}_t), u_t \hat{u}_t \rangle \leq 0 \quad for \ all \ t \in \{0, ..., h\}, \ for \end{array}$ all $u_t \in U_t$.
- (4) For all $t \in \{1, ..., h + 1\}$, there exists $a_t, b_t \in \mathbb{R}_+$ such that, for all $s \in \mathbb{R}_+$ $\{1, ..., h\}, \|p_t^h\| \le a_t \lambda_0^h + b_t \|p_s^h\|.$
- (5) $(\lambda_0^h, p_{1|Z_0}^h, p_{2|Z_1}^h) \neq (0, 0, 0).$
- (6) For all $z_0 \in Z_0$, for all $z_1 \in Z_1$, there exists $c_{z_0,z_1} \in \mathbb{R}$ such that, for all $h \ge 2$, $p_1^h(z_0) + p_2^h(z_1) \le c_{z_0, z_1} \lambda_0^h$.
- (7) For all $v \in X$ there exists $(z_0, z_1) \in Z_0 \times Z_1$ such that $p_2^h(v) = p_1^h(z_0) + p_2^h(z_1) \lambda_0^h D_1 \phi_1(\hat{x}_1, \hat{u}_1)(z_0)$ for all $h \ge 2$.

Proof. Proof of (1)-(4) Note that conditions (4.3) and (4.4) are consequences of (H4). We use λ_0^h and $(p_{t+1}^h)_{0 \le t \le h}$ which are provided by Lemma 4.3. Hence conclusions (1), (2) and (3) are given by Lemma 4.3. The conclusion (4) is the conclusion (e) of Lemma 4.3.

Proof of (5) From the conclusion (f) of Lemma 4.3, we know that $(\lambda_0^h, p_1^h) \neq (0, 0)$. We want to prove that $[(\lambda_0^h, p_1^h) \neq (0, 0)]$ implies (5). To do that we proceed by contraposition; we assume that $[\lambda_0^h = 0, p_1^h|_{Z_0} = 0, p_2^h|_{Z_1} = 0]$ and we want to prove that $[\lambda_0^h = 0, p_1^h = 0]$. Since $\lambda_0^h = 0$, using the conclusion (2) we obtain $p_1^h = p_2^h \circ D_1 f_1(\hat{x}_1, \hat{u}_1)$ which implies

$$p_1^h \circ D_2 f_0(\sigma, \hat{u}_0)(T_{U_0}(\hat{u}_0)) = p_2^h \circ D_1 f_1(\hat{x}_1, \hat{u}_1) \circ D_2 f_0(\sigma, \hat{u}_0)(T_{U_0}(\hat{u}_0)),$$

and since $p_{1|Z_0}^h = 0$, we obtain $p_2^h \circ D_1 f_1(\hat{x}_1, \hat{u}_1) \circ D_2 f_0(\sigma, \hat{u}_0)(T_{U_0}(\hat{u}_0)) = 0$, and since $p_{2|Z_1}^h = 0$, using (H5), we obtain $p_2^h = 0$ (on X all over). Hence $p_1^h = 0$ $p_2^h \circ D_1 f_1(\hat{x}_1, \hat{u}_1) = 0$. The proof of (5) is complete.

Proof of (6) Let $z_0 \in Z_0$, $z_1 \in Z_1$. Using conclusion (g) of Lemma 4.3, we obtain that there exists $c_{z_0}^0 \in \mathbb{R}$ such that $p_1^h(z_0) \leq c_{z_0}^0 \lambda_0^h$ for all $h \geq 1$, and that there exists $c_{z_1}^1 \in \mathbb{R}$ such that $p_2^h(z_1) \leq c_{z_1}^1 \lambda_0^h$ for all $h \geq 2$. Setting $c_{z_0,z_1} := c_{z_0}^0 + c_{z_1}^1$ we obtain the announced conclusion.

Proof of (7) From (H5), for all $v \in X$, there exist $\zeta_0 \in T_{U_0}(\hat{u}_0)$ and $\zeta_1 \in T_{U_1}(\hat{u}_1)$ such that

$$v = D_1 f_1(\hat{x}_1, \hat{u}_1) \circ D_2 f_0(\sigma, \hat{u}_0)(\zeta_0) + D_2 f_1(\hat{x}_1, \hat{u}_1)(\zeta_1).$$

We set $z_0 := D_2 f_0(\sigma, \hat{u}_0)(\zeta_0) \in Z_0$ and $z_1 := D_2 f_1(\hat{x}_1, \hat{u}_1)(\zeta_1) \in Z_1$, hence we have

(4.28)
$$v = D_1 f_1(\hat{x}_1, \hat{u}_1)(z_0) + z_1.$$

From conclusion (2) we deduce

$$p_1^h \circ D_2 f_0(\sigma, \hat{u}_0) = p_2^h \circ D_1 f_1(\hat{x}_1, \hat{u}_1) \circ D_2 f_0(\sigma, \hat{u}_0) + \lambda_0^h D_1 \phi_1(\hat{x}_1, \hat{u}_1) \circ D_2 f_0(\sigma, \hat{u}_0).$$

Applying this last equation to ζ_0 we obtain

$$p_1^h(z_0) = p_2^h \circ D_1 f_1(\hat{x}_1, \hat{u}_1)(z_0) + \lambda_0^h D_1 \phi_1(\hat{x}_1, \hat{u}_1)(z_0).$$

M. BACHIR AND J. BLOT

Adding $p_2^h(z_1)$ to this equality we obtain

 $p_1^h(z_0) + p_2^h(z_1) = p_2^h \circ D_1 f_1(\hat{x}_1, \hat{u}_1)(z_0) + p_2^h(z_1) + \lambda_0^h D_1 \phi_1(\hat{x}_1, \hat{u}_1)(z_0).$

Using (4.28) we have $p_1^h(z_0) + p_2^h(z_1) = p_2^h(v) + \lambda_0^h D_1 \phi_1(\hat{x}_1, \hat{u}_1)(z_0)$ which implies the announced equality.

5. Proof of the main theorem

Proposition 4.5 provides sequences $(\lambda_0^h)_{h\geq 2}$, $(p_t^h)_{h\geq t}$ for all $t\in \mathbb{N}_*$. We set $q_1^h:=p_1^h\circ D_2f_0(\sigma,\hat{u}_0)\in U^*$ and $q_2^h:=p_2^h\circ D_2f_1(\hat{x}_1,\hat{u}_1)\in U^*$ for all $h\geq 2$. From conclusion (5) of Proposition 4.5 we obtain

$$(\lambda_0^h, q_1^h|_{T_{U_0}(\hat{u}_0)}, q_2^h|_{T_{U_1}(\hat{u}_1)}) \neq (0, 0, 0).$$

We introduce $\Sigma := \overline{\operatorname{aff}}(T_{U_0}(\hat{u}_0) \times T_{U_1}(\hat{u}_1))$ the closed affine hull of $T_{U_0}(\hat{u}_0) \times T_{U_1}(\hat{u}_1)$ which is a closed vector subspace since the tangent cones contain the origin. From the previous relation we can assert that $(\lambda_0^h, (q_1^h, q_2^h)|_{\Sigma}) \neq (0, (0, 0))$. We introduce the number

$$\theta^h := \lambda_0^h + \|(q_1^h, q_2^h)|_{\Sigma}\|_{\Sigma^*} > 0.$$

Since the list of the multipliers of the problem in finite horizon is a cone, we can replace λ_0^h by $\frac{1}{\theta^h} \lambda_0^h$ and the p_t^h by $\frac{1}{\theta^h} p_t^h$ (without to change the writing), and so we can assume that the following property holds.

(5.1)
$$\forall h \ge 2, \ \lambda_0^h + \|(q_1^h, q_2^h)|_{\Sigma}\|_{\Sigma^*} = 1.$$

Using the Banach-Alaoglu theorem, we can assert that there exist an increasing mapping $\varphi_1 : [2, +\infty)_{\mathbb{N}} \to [2, +\infty)_{\mathbb{N}}, \lambda_0 \in \mathbb{R}, (q_1, q_2) \in \Sigma^*$ such

$$(\lambda_0^{\varphi_1(h)}, (q_1^{\varphi_1(h)}, q_2^{\varphi_1(h)})|_{\Sigma}) \xrightarrow{w^*} (\lambda_0, (q_1, q_2)) \text{ when } h \to +\infty.$$

Now we want to establish that

(5.2)
$$(\lambda_0, (q_1, q_2)) \neq (0, (0, 0)).$$

To do that we proceed by contradiction; we assume that $\lambda_0 = 0$ and $(q_1, q_2) = (0, 0)$. From conclusion (6) of Proposition 4.5 we deduce that, for all $\zeta_0 \in T_{U_0}(\hat{u}_0)$ and for all $\zeta_1 \in T_{U_1}(\hat{u}_1)$, there exists $c_{\zeta_0,\zeta_1} \in \mathbb{R}$ such that $q_1^{\varphi_1(h)}(\zeta_0) + q_2^{\varphi_1(h)}(\zeta_1) \leq c_{\zeta_0,\zeta_1}\lambda_0^{\varphi_1(h)}$ for all $h \geq 2$. Hence we can use Proposition 3.2 with $Y = \Sigma$, $K = T_{U_0}(\hat{u}_0) \times T_{U_1}(\hat{u}_1)$, $S = \Sigma$, $\rho_h = \lambda_0^{\varphi_1(h)}$, and $\pi_h = (q_1^{\varphi_1(h)}, q_2^{\varphi_1(h)})_{|\Sigma}$. Consequently we obtain that $\lim_{h \to +\infty} \|(q_1^{\varphi_1(h)}, q_2^{\varphi_1(h)})|_{\Sigma}\|_{\Sigma^*} = 0$. Since we also have $\lim_{h \to +\infty} \lambda_0^{\varphi_1(h)} = 0$, we obtain a contradiction with (5.1). Hence (5.2) is proven.

From conclusion (7) of Proposition 4.5 we have, for all $x \in X$, there exists $(\zeta_0, \zeta_1) \in \Sigma$ such that, for all $h \ge 2$,

$$p_2^{\varphi_1(h)}(x) = (q_1^{\varphi_1(h)}, q_2^{\varphi_1(h)})_{|\Sigma|}(\zeta_0, \zeta_1) - \lambda_0^{\varphi_1(h)} D_1 \phi_1(\hat{x}_1, \hat{u}_1) \circ D_2 f_0(\sigma, \hat{u}_0)(\zeta_0)$$

which permits to say that there exists $p_2 \in X^*$ such that $p_2^{\varphi_1(h)} \xrightarrow{w^*} p_2$ when $h \to +\infty$. From conclusion (2) of Proposition 4.5 at t = 1, we obtain that there exists $p_1 \in X^*$ such that $p_1^{\varphi_1(h)} \xrightarrow{w^*} p_1$ when $h \to +\infty$, and from (5.2) we obtain

(5.3)
$$(\lambda_0, (p_1, p_2)) \neq (0, (0, 0)).$$

32

Since $(p_1^{\varphi_1(h)})_{h\geq 2}$ is weak-star convergent on X, using the Banach-Steinhaus theorem we can assert that the sequence $(||p_1^{\varphi_1(h)}||_{X^*})_{h\geq 2}$ is bounded. Since $(\lambda_0^{\varphi_1(h)})_{h\geq 2}$ is convergent in \mathbb{R} , it is bounded, and from conclusion (4) of Proposition 4.5, we deduce that the sequence $(p_t^{\varphi_1(h)})_{h\in[t,+\infty)_{\mathbb{N}}}$ is bounded for each $t \in \mathbb{N}_*$. Then we can use Proposition 3.3 to ensure the existence of an increasing function $\varphi_2 : [2,+\infty)_{\mathbb{N}} \to$ $[2,+\infty)_{\mathbb{N}}$ and a sequence $(p_t)_{t\in N_*} \in (X^*)^{\mathbb{N}_*}$ such that $p_t^{\varphi_1\circ\varphi_2(h)} \stackrel{w^*}{\to} p_t$ when $h \to +\infty$ for all $t \in \mathbb{N}_*$, and we have also $\lambda_0^{\varphi_1\circ\varphi_2(h)} \to \lambda_0$ when $h \to +\infty$. Hence we have built all the multipliers. The properties of these multipliers are obtained by taking limits from the properties of the λ_0^h and the p_t^h . Their non triviality is proven by (5.3). Hence the proof of the main result is complete.

References

- [1] J.-P. Aubin and I. Ekeland, Applied nonlinear analysis, John Wiley & Sons, New York, 1984.
- [2] M. Bachir and J. Blot Infinite Dimensional Infinite-horizon Pontryagin principles for discretetime problems, Set-Valued Var. Anal. 23 (2015) 43-54.
- [3] M. Bachir and J. Blot Infinite dimensional multipliers and Pontryagin principles for discretetime problems, Pure Applied Func. Anal. 2 (2017), 411426.
- [4] J. Blot and P. Cieutat, Completness of sums of subspaces of bounded functions and applications, Commun. Math. Anal. 19(2) (2018), 43-61.
- [5] J. Blot and H. Chebbi, Discrete time Pontryagin principle in infinite horizon, J. Math. Anal. Appl. 246 (2000), 265-279.
- [6] J. Blot and N. Hayek, Infinite-horizon optimal control in the discrete-time framework, Springer, New York, 2014.
- [7] D.A. Carlson, A.B. Haurie and A. Leizarowitz, Infinite horizon optimal control; deterministic and stochastic systems, Second revised edition, Springer, Berlin, 1991.
- [8] W.C. Clark, Mathematical bioeconomics; optimal management of renewable resources, second edition, John Wiley and Sons, Inc;, Hoboken, New Jersey, 2005.
- [9] J. Jahn, Introduction to the theory of nonlinear optimization, Third edition, Springer-Verlag, Berlin, 2007.
- [10] P. Michel, Some clarifications on the transversality condition, Econometrica 58 (1990), 705-728.
- [11] N. L. Stokey, R.E. Lucas and E.C. Prescott, Recursive methods in economic dynamics, Harvard University Press, Cambridge, MA, 1989.
- [12] C. Zălinescu, Convex analysis in general vector spaves, World Scientific Publishing Co., Pte, Ltd, Singapore, 2002.
- [13] A.J. Zaslavski, Turnpike properties in the calculus of variations and optimal Control, Springer Science+Business Media, New york, N.Y., 2006.

Manuscript received June 26 2017 revised July 25 2017

M. BACHIR

Laboratoire SAMM EA4543, Université Paris 1 Panthéon-Sorbonne, centre P.M.F., 90 rue de Tolbiac, 75634 Paris cedex 13, France

E-mail address: Mohammed.Bachir@univ-paris1.fr

J. Blot

Laboratoire SAMM EA4543, Université Paris 1 Panthéon-Sorbonne, centre P.M.F., 90 rue de Tolbiac, 75634 Paris cedex 13, France

E-mail address: blot@univ-paris1.fr