

# MULTIPLE PERIODIC SOLUTIONS OF INFINITE-DIMENSIONAL PENDULUM-LIKE EQUATIONS

ALESSANDRO FONDA, JEAN MAWHIN, AND MICHEL WILLEM

ABSTRACT. We prove the multiplicity of periodic solutions for an equation in a separable Hilbert space H, with T-periodic dependence in time, of the type

$$\ddot{x} + \mathcal{A}x + \nabla_x V(t, x) = e(t) \,.$$

Here,  $\mathcal{A}$  is a semi-negative definite bounded selfadjoint operator, with nontrivial null-space  $\mathcal{N}(\mathcal{A})$ , the function V(t, x) is bounded above, periodic in x along a basis of  $\mathcal{N}(\mathcal{A})$ , with  $\nabla_x V$  having its image in a compact set, and e(t) has mean value in  $\mathcal{N}(\mathcal{A})^{\perp}$ . Our results generalize several well-known theorems in the finite-dimensional setting, as well as a recent existence result in [1].

## 1. INTRODUCTION

Motivated by the model of a periodically forced pendulum, the existence of at least two geometrically distinct T-periodic solutions for a scalar differential equation of the form

$$\ddot{x} + \partial_x V(t, x) = e(t)$$

was first proved in [16], using the direct method of the calculus of variations and the Mountain Pass Theorem, assuming V(t, x) to be *T*-periodic with respect to *t* and  $\tau$ -periodic with respect to *x*, and e(t) to be *T*-periodic with zero mean, i.e.,

(1.1) 
$$\int_{0}^{T} e(t) dt = 0$$

This result extended an existence theorem first proved in [12], and later rediscovered independently in [5, 21].

Here, and in the sequel, for simplicity all functions will be assumed to be continuous. It can be seen that the multiplicity result in [16] is optimal if no further conditions are added. Different proofs have also been provided, e.g., in [8, 9, 11], by the use of some generalized versions of the Poincaré–Birkhoff theorem.

The result in [16] was later generalized in [17], through a similar approach, to the corresponding system in  $\mathbb{R}^N$ ,

(1.2) 
$$\ddot{x} + \nabla_x V(t, x) = e(t),$$

<sup>2010</sup> Mathematics Subject Classification. 34C25, 34G20, 47J30.

Key words and phrases. pendulum equation, periodic solutions, BVP in Hilbert space.

when  $V(t, x) = V(t, x_1, \ldots, x_N)$  is *T*-periodic in *t* and  $\tau_k$ -periodic in  $x_k$ , for every  $k = 1, \ldots, N$ . The first aim of this paper is to extend such a result to an infinitedimensional setting. So, let *H* be a separable Hilbert space, and let  $(e_k)_{k\geq 1}$  be a Hilbert basis. We assume  $V : \mathbb{R} \times H \to \mathbb{R}$  to be continuous, *T*-periodic with respect to its first variable *t*, and continuously differentiable with respect to its second variable *x*. Here is our result.

**Theorem 1.1.** Assume that there exists a sequence of positive real numbers  $(\tau_k)_{k\geq 1}$  such that

(1.3) 
$$V(t, x + \tau_k e_k) = V(t, x), \text{ for every } (t, x) \in [0, T] \times H$$
and  $k = 1, 2, \dots$ 

If  $\sum_{k=1}^{\infty} \tau_k^2 < +\infty$ , then equation (1.2) has at least two geometrically distinct *T*-periodic solutions, for every e(t) satisfying (1.1).

The above theorem generalizes [1, Theorem 6], where further regularity assumptions were made on V, in order to obtain the existence of *at least one* T-periodic solution. A Galerkin-type argument was used there to reduce the problem to a sequence of finite-dimensional differential systems, to which a generalized version of the Poincaré–Birkhoff theorem applies (cf. [9]), followed by a limit process.

The proof of Theorem 1.1 will follow the same ideas introduced in [16, 17], taking advantage of the compactness of the Hilbert cube  $\prod_{k=1}^{\infty} [0, \tau_k]$ . The first solution will be obtained by minimization of the action functional, while the second one will be of mountain pass type.

Using the Lusternik–Schnirelmann theory, it was proved in [15] that, under the same assumptions, system (1.2) in  $\mathbb{R}^N$  has indeed at least N+1 geometrically distinct T-periodic solutions, thus generalizing the result in [17]. (Notice that, when  $N \geq 2$ , the multiplicity result is not optimal, as shown by the four equilibria of a double pendulum.) Even more, a system of the type

(1.4) 
$$\ddot{x} + \mathcal{A}x + \nabla_x V(t, x) = e(t)$$

was considered there, involving a symmetric matrix  $\mathcal{A}$ . Other results in this direction, including the case of Hamiltonian systems leading to a strongly indefinite action functional, were studied, e.g., in [3, 4, 6, 7, 9, 10, 13, 14, 20].

The second aim of this paper is to obtain multiplicity results for an infinitedimensional system of the type (1.4), when  $\mathcal{A} : H \to H$  is a semi-negative definite bounded selfadjoint operator, whose spectrum contains 0 as an isolated eigenvalue, V(t,x) is bounded above and *T*-periodic in *t*, and the image of  $\nabla_x V$  is contained in a compact set of *H*. Denoting by  $\mathcal{N}(\mathcal{A})$  the null space of  $\mathcal{A}$ , we distinguish two cases.

If  $\mathcal{N}(\mathcal{A})$  has finite dimension N and V(t, x) satisfies a periodicity condition of the type (1.3), with the  $e_k$  replaced by the elements of an orthonormal basis of  $\mathcal{N}(\mathcal{A})$ , the existence of at least N + 1 geometrically distinct T-periodic solutions is proved, when the mean value of e(t) belongs to  $\mathcal{N}(\mathcal{A})^{\perp}$ . The precise statement will be given in Section 2. The proof, provided in Section 3, will be carried out by the use of an abstract theorem, given in [18] and inspired by [19], providing the multiplicity of

critical points of some functionals in a Banach space X which are bounded below, invariant under the action of some discrete subgroups of X, and satisfy a suitable Palais–Smale condition.

If  $\mathcal{N}(\mathcal{A})$  has infinite dimension, assuming in addition that  $\sum_{k=1}^{\infty} \tau_k^2 < +\infty$ , after finding the first solution by minimization of the action functional, a second one is provided by the Mountain Pass Theorem. We thus get, in this case, the existence of *at least two* geometrically distinct *T*-periodic solutions.

The paper ends with some examples and an open problem.

### 2. The main result

Let *H* be a separable Hilbert space, with scalar product  $(\cdot, \cdot)$  and corresponding norm  $|\cdot|$ . In this space, we consider the equation

(2.1) 
$$\ddot{x} + \mathcal{A}x + \nabla_x V(t, x) = e(t),$$

where  $\mathcal{A} \in \mathcal{L}(H)$  is a bounded selfadjoint operator, and  $e : \mathbb{R} \to H$  is continuous and T-periodic. Concerning the function  $V : \mathbb{R} \times H \to \mathbb{R}$ , it is continuous, T-periodic in its first variable t, and differentiable with respect to its second variable x, with corresponding continuous gradient  $\nabla_x V : \mathbb{R} \times H \to H$ .

Let us introduce our assumptions. We denote by  $\mathcal{N}(\mathcal{A})$  the null-space of  $\mathcal{A}$ , and by  $\sigma(\mathcal{A})$  its spectrum. We take a Hilbert basis  $(a_k)_k$  of  $\mathcal{N}(\mathcal{A})$ , considered as a subspace of H. If  $\mathcal{N}(\mathcal{A})$  has a finite dimension, its basis will be given by  $(a_1, \ldots, a_N)$ ; if it is infinite-dimensional, we will have a sequence of vectors  $(a_1, a_2, \ldots)$ .

**A1.** The selfadjoint operator  $\mathcal{A}$  is semi-negative definite, with  $\mathcal{N}(\mathcal{A}) \neq \{0\}$ , and

$$\sup\left(\sigma(\mathcal{A})\setminus\{0\}\right)<0.$$

So, 0 is an isolated point of  $\sigma(\mathcal{A})$ .

**A2.** The mean value of e(t) is orthogonal to  $\mathcal{N}(\mathcal{A})$ , i.e.,

$$\int_0^T e(t) \, dt \in \mathcal{N}(\mathcal{A})^\perp \, .$$

Then, we have that

$$\int_0^T (e(t), a_k) \, dt = 0 \,, \text{ for every } k = 1, 2, \dots$$

**A3.** There exists a sequence of positive real numbers  $(\tau_k)_{k\geq 1}$  such that

 $V(t, x + \tau_k a_k) = V(t, x)$ , for every  $(t, x) \in [0, T] \times H$  and  $k = 1, 2, \dots$ 

A4. There is a nonnegative constant C such that

$$V(t, x) \leq C$$
, for every  $(t, x) \in [0, T] \times H$ .

A5. The set  $\nabla_x V([0,T] \times H)$  is precompact in H.

In the above setting, we can now state the main result of this paper.

**Theorem 2.1.** Assume that conditions A1 to A5 hold. If  $\mathcal{N}(\mathcal{A})$  is finite-dimensional, then equation (2.1) has at least dim  $\mathcal{N}(\mathcal{A}) + 1$  geometrically distinct *T*-periodic solutions. On the other hand, if  $\mathcal{N}(\mathcal{A})$  is infinite-dimensional and  $\sum_{k=1}^{\infty} \tau_k^2 < +\infty$ , then there are at least two of them.

Notice that, once a *T*-periodic solution x(t) has been found, any function obtained by adding to it some integer multiples of  $\tau_k a_k$  is still a *T*-periodic solution. We say that two *T*-periodic solutions are *geometrically distinct* if they cannot be obtained one from the other in this way.

Concerning assumption A5, we remark that it will surely be satisfied if the following holds.

**A5'.** There exists a Hilbert basis  $(e_k)_{k\geq 1}$  of H and a nonnegative sequence  $(M_k)_k$ , with  $\sum_{k=1}^{\infty} M_k^2 < +\infty$ , such that

$$\left|\frac{\partial V}{\partial e_k}(t,x)\right| \le M_k$$
, for every  $(t,x) \in [0,T] \times H$  and  $k = 1, 2, ...$ 

Indeed, A5' implies that  $\nabla_x V([0,T] \times H)$  is contained in a Hilbert cube, which is a compact set in H. In the above formula, we have used the notation

$$\frac{\partial V}{\partial e_k}(t,x) = \lim_{\tau \to 0} \frac{V(t,x+\tau e_k) - V(t,x)}{\tau}$$

Notice that Theorem 1.1 is a direct consequence of Theorem 2.1, taking  $\mathcal{A} = 0$  and  $(a_k)_k = (e_k)_k$ , a Hilbert basis of H. Indeed, the periodicity assumption in Theorem 1.1 and the compactness of the set  $[0,T] \times \prod_{k=1}^{\infty} [0,\tau_k]$  show that A4 and A5 are surely satisfied.

In the proof of Theorem 2.1, we will need a result from [18], which we now recall, for the reader's convenience.

Let G be a discrete subgroup of a Banach space X and  $\pi : X \to X/G$  be the canonical surjection. A subset S of X is G-invariant if  $\pi^{-1}(\pi(S)) = S$ , and a function f defined on X is G-invariant if f(u+g) = f(u), for every  $u \in X$  and every  $g \in G$ . If  $\varphi \in C^1(X, \mathbb{R})$  is G-invariant, then  $\varphi'$  is also G-invariant, and if u is a critical point of  $\varphi$ , the same is true for u + g for all  $g \in G$ . The corresponding set  $\{u+g: g \in G\}$  is called a critical orbit of  $\varphi$ .

A G-invariant differentiable function  $\varphi : X \to \mathbb{R}$  satisfies the  $(PS)_G$  condition if, for every sequence  $(u_n)_n$  in X such that  $\varphi(u_n)$  is bounded and  $\varphi'(u_n) \to 0$ , the sequence  $(\pi(u_n))_n$  contains a convergent subsequence.

The following multiplicity result for the critical points of G-invariant functionals is stated as Theorem 4.12 in [18].

**Theorem 2.2.** Let  $\varphi \in C^1(X, \mathbb{R})$  be a *G*-invariant functional satisfying the  $(PS)_G$  condition. If  $\varphi$  is bounded from below and if the dimension N of the space generated by G is finite, then  $\varphi$  has at least N + 1 critical orbits.

### 3. Proof of Theorem 2.1

Let  $L^2([0,T], H)$  be the space of measurable functions  $x : [0,T] \to H$  such that |x| is square integrable. It is a Hilbert space equipped with the scalar product

$$\langle x, y \rangle_2 = \int_0^T (x(t), y(t)) dt$$

and corresponding norm

$$||x||_2 = \left(\int_0^T |x(t)|^2 dt\right)^{\frac{1}{2}}.$$

We consider the space  $H^1([0,T], H)$ , made of those functions x belonging to  $L^2([0,T], H)$  with weak derivative  $\dot{x}$  also in  $L^2([0,T], H)$ . It is a Hilbert space, as well, with the scalar product

$$\langle x, y \rangle = \langle x, y \rangle_2 + \langle \dot{x}, \dot{y} \rangle_2 = \int_0^T \left[ (x(t), y(t)) + (\dot{x}(t), \dot{y}(t)) \right] dt \,,$$

and corresponding norm

$$||x|| = \left(||x||_{2}^{2} + ||\dot{x}||_{2}^{2}\right)^{\frac{1}{2}} = \left(\int_{0}^{T} \left[|x(t)|^{2} + |\dot{x}(t)|^{2}\right] dt\right)^{\frac{1}{2}}.$$

Moreover,  $H^1([0,T], H)$  is continuously embedded in C([0,T], H), the space of continuous functions, with the usual norm

$$||x||_{\infty} = \max\{|x(t)| : t \in [0,T]\}.$$

(For further information on the space  $H^1([0,T], H)$  we refer, e.g., to [2].)

Let

$$H_T^1 = \left\{ x \in H^1([0,T],H) : x(0) = x(T) \right\},\$$

and define the functional  $\varphi: H^1_T \to \mathbb{R}$  as

$$\varphi(x) = \int_0^T \left[ \frac{1}{2} |\dot{x}(t)|^2 - \frac{1}{2} (\mathcal{A}x(t), x(t)) - V(t, x(t)) + (e(t), x(t)) \right] dt \,.$$

It is continuously differentiable, and its critical points correspond to the T-periodic solutions of (2.1). Moreover, by A2 and A3,

(3.1) 
$$\varphi(x + \tau_k a_k) = \varphi(x)$$
, for every  $x \in H^1_T$  and  $k \ge 1$ .

As usual, we identify the constant functions with their constant value. So, having identified H with the space of constant functions, it will be a subspace of  $H_T^1$ . Hence, we can write

$$H^1_T = H \oplus W = \mathcal{N}(\mathcal{A}) \oplus \mathcal{N}(\mathcal{A})^{\perp} \oplus W = \mathcal{N}(\mathcal{A}) \oplus W.$$

Here, W is the orthogonal space to H in  $H_T^1$ ,  $\mathcal{N}(\mathcal{A})^{\perp}$  is the orthogonal to  $\mathcal{N}(\mathcal{A})$ in H, and  $\widetilde{W} = \mathcal{N}(\mathcal{A})^{\perp} \oplus W$ . Correspondingly, we will write each  $x \in H_T^1$  as  $x(t) = \bar{x} + \tilde{x}(t)$ , with  $\bar{x} \in \mathcal{N}(\mathcal{A})$  and  $\tilde{x} \in \widetilde{W}$ . Moreover, we will write  $\tilde{x}(t) = \hat{x} + \check{x}(t)$ , with  $\hat{x} \in \mathcal{N}(\mathcal{A})^{\perp}$  and  $\check{x} \in W$ . Notice that, for any  $x \in H_T^1$ ,

(3.2) 
$$[x] := \frac{1}{T} \int_0^T x(t) \, dt = \bar{x} + \hat{x} \,, \quad \frac{1}{T} \int_0^T \check{x}(t) \, dt = 0 \,.$$

**Proposition 3.1.** For every  $x \in H_T^1$ , one has

$$\|\check{x}\|_{\infty} \le \sqrt{T} \, \|\dot{x}\|_2$$

*Proof.* Let  $(e_k)_{k\geq 1}$  be a Hilbert basis of H. Then, for any function  $x \in H^1_T$ , we may write

$$\check{x}(t) = \sum_{k=1}^{\infty} (\check{x}(t), e_k) e_k = \sum_{k=1}^{\infty} \check{x}_k(t) e_k.$$

Being  $\check{x}_k$  continuous, T-periodic with zero mean, there is a  $t_k \in [0,T]$  for which  $\check{x}_k(t_k) = 0$ , hence

$$|\check{x}_k(t)| = \left|\check{x}_k(t_k) + \int_{t_k}^t \dot{x}_k(s) \, ds\right| \le \int_0^T |\dot{x}_k(s)| \, ds \le \sqrt{T} \left(\int_0^T |\dot{x}_k(s)|^2 \, ds\right)^{\frac{1}{2}},$$

for every  $t \in [0, T]$ . As a consequence,

$$|\check{x}(t)|^2 = \sum_{k=1}^{\infty} |\check{x}_k(t)|^2 \le T \int_0^T \sum_{k=1}^{\infty} |\dot{x}_k(s)|^2 \, ds = T \int_0^T |\dot{x}(s)|^2 \, ds$$

for every  $t \in [0, T]$ , whence the conclusion.

By A1, A2, A4, (3.2) and (3.3), setting  $\delta := -\sup(\sigma(A) \setminus \{0\}),$  $\varphi(x) = \int_0^T \left[ \frac{1}{2} |\dot{x}(t)|^2 - \frac{1}{2} (\mathcal{A}\tilde{x}(t), \tilde{x}(t)) - V(t, x(t)) + (e(t), \tilde{x}(t)) \right] dt$ 

$$\geq \int_{0} \left[ \frac{1}{2} |\dot{x}(t)|^{2} - \frac{1}{2} (\mathcal{A}\hat{x}, \hat{x}) - \frac{1}{2} (\mathcal{A}\check{x}(t), \check{x}(t)) \right] dt - CT - T \|e\|_{\infty} \|\tilde{x}\|_{\infty}$$

$$\geq \int_{0}^{T} \frac{1}{2} |\dot{x}(t)|^{2} dt - \frac{1}{2} T (\mathcal{A}\hat{x}, \hat{x}) - CT - T \|e\|_{\infty} (|\hat{x}| + \|\check{x}\|_{\infty})$$

$$\geq \frac{1}{2} \|\dot{x}\|_{2}^{2} + \frac{1}{2} T \delta |\hat{x}|^{2} - CT - T^{\frac{3}{2}} \|e\|_{\infty} \|\dot{x}\|_{2} - T \|e\|_{\infty} |\hat{x}| .$$

Hence, since  $\delta > 0$ , there are two positive constants c > 0 and c' > 0 for which  $\varphi(x) \ge c \left( \|\dot{x}\|_2^2 + |\hat{x}|^2 \right) - c',$ (3.4)

and the functional  $\varphi$  is bounded below.

For  $u \in C([0,T], H)$ , we denote by Pu the indefinite integral defined on [0,T] by

$$Pu(t) = \int_0^t u(s) \, ds \, .$$

**Lemma 3.2.** Let  $E \subseteq C([0,T],H)$  be such that  $A := \{u([0,T]) : u \in E\}$  is precompact in H. Then:

- (a) the set  $B := \{\int_0^T u(t) dt : u \in E\}$  is precompact in H; (b) the set  $S := \{Pu : u \in E\}$  is precompact in C([0,T], H).

956

*Proof.* (a) Let  $\varepsilon > 0$ . There exists a finite sequence  $(v_1, \ldots, v_n)$  in H such that, denoting by  $B(u, \rho)$  any open ball of center u and radius  $\rho$ ,

$$A \subseteq \bigcup_{k=1}^n B(v_k, \varepsilon)$$

We denote by  $Q_0$  the orthogonal projection from H to the space V generated by  $(v_1, \ldots, v_n)$ . The set

$$C = \left\{ \int_0^T Q_0 u(t) \, dt : u \in E \right\}$$

is bounded in V, hence precompact in V. This implies the existence of a finite sequence  $(w_1, \ldots, w_m)$  in V such that

$$C \subseteq \bigcup_{k=1}^m B(w_k, \varepsilon) \,.$$

For every  $u \in E$ , we have

$$\left|\int_0^T u(t) \, dt - \int_0^T Q_0 u(t) \, dt\right| \le \int_0^T |u(t) - Q_0 u(t)| \, dt \le \varepsilon T \, .$$

It follows that

$$B \subseteq \bigcup_{k=1}^{m} B(w_k, (1+T)\varepsilon)$$

Since  $\varepsilon$  is arbitrary, B is precompact in H.

(b) Let us define

$$R := \{ P(Q_0 u) : u \in E \} \,.$$

The set  $\{P(Q_0u)(t) : t \in [0,T], u \in E\}$  is bounded in V, hence precompact in V. For  $0 \le t_1 \le t_2 \le T$ , we have

$$|P(Q_0u)(t_2) - P(Q_0u)(t_1)| = \left| \int_{t_1}^{t_2} Q_0(u)(s) \, ds \right| \le c(t_2 - t_1) \,,$$

for some c > 0. By the Ascoli–Arzelá theorem, the set R is precompact in C([0, T], V). This implies, for any  $\varepsilon > 0$ , the existence of a finite sequence  $(f_1, \ldots, f_N)$  in C([0, T], V) such that

$$R \subseteq \bigcup_{k=1}^N B(f_k, \varepsilon) \,.$$

Since, for every  $u \in E$  and  $t \in [0, T]$ , we have

$$|Pu(t) - P(Q_0 u)(t)| \le \int_0^t |u(s) - Q_0 u(s)| \, ds \le T\varepsilon \,,$$

we conclude that

$$S \subseteq \bigcup_{k=1}^{N} B(f_k, (1+T)\varepsilon).$$

Since  $\varepsilon > 0$  is arbitrary, S is precompact in C([0, T], H).

We now prove the following.

**Proposition 3.3.** If  $(x^n)_n$  is a sequence in  $H^1_T$  such that  $(\varphi(x^n))_n$  is bounded and  $\nabla \varphi(x^n) \to 0$ , then  $(\tilde{x}^n)_n$  has a convergent subsequence.

Proof. Since  $(\varphi(x^n))_n$  is bounded, by (3.3) and (3.4) we have that  $(\tilde{x}^n)_n$  is bounded in  $H_T^1$ . On the other hand, we can modify  $\bar{x}^n$  into some  $\bar{z}^n$  such that the scalar product  $(\bar{z}^n, a_k)$  belongs to  $[0, \tau_k]$ , for every k, and  $(\bar{z}^n, a_k) \equiv (\bar{x}^n, a_k) \mod \tau_k$ . Defining  $z^n = \bar{z}^n + \tilde{x}^n$ , we have a new sequence for which  $\varphi(z^n) = \varphi(x^n)$  and  $\nabla \varphi(z^n) = \nabla \varphi(x^n)$ , by (3.1). Moreover,  $(z^n)_n$  is bounded, hence there is a subsequence, still denoted by  $(z^n)_n$ , which weakly converges to some  $z^* \in H_T^1$ . We want to show that  $(z^n)_n$  strongly converges to  $z^*$  in  $H_T^1$ .

Since  $\nabla \varphi(z^n) \to 0$  and  $(z^n)_n$  weakly converges to  $z^*$ , we have that

$$\langle \nabla \varphi(z^n) - \nabla \varphi(z^*), z^n - z^* \rangle \to 0,$$

i.e.,

(3.5) 
$$\lim_{n} \int_{0}^{T} \left[ |\dot{z}^{n}(t) - \dot{z}^{*}(t)|^{2} - (\mathcal{A}(z^{n}(t) - z^{*}(t)), z^{n}(t) - z^{*}(t)) - (\nabla_{x}V(t, z^{n}(t)) - \nabla_{x}V(t, z^{*}(t)), z^{n}(t) - z^{*}(t)) \right] dt = 0$$

Since  $(z^n)_n$  weakly converges to  $z^*$  in  $L^2([0,T],H)$ ,

(3.6) 
$$\lim_{n} \int_{0}^{1} \left( \nabla_{x} V(t, z^{*}(t)), z^{n}(t) - z^{*}(t) \right) dt = 0.$$

Claim. Up to a subsequence,

(3.7) 
$$\lim_{n} \int_{0}^{T} (\nabla_{x} V(t, z^{n}(t)), z^{n}(t) - z^{*}(t)) dt = 0.$$

*Proof of the Claim.* Define on [0, T] the continuous functions

$$w^{n}(t) = \nabla_{x} V(t, z^{n}(t)), \quad y^{n}(t) = z^{n}(t) - z^{*}(t),$$

having values in H. Using the notation in (3.2), we have

$$\int_0^T (w^n(t), y^n(t)) dt = \int_0^T \left( [w^n] + \check{w}^n(t), [y^n] + \check{y}^n(t) \right) dt$$
$$= T([w^n], [y^n]) + \int_0^T (\check{w}^n(t), \check{y}^n(t)) dt.$$

Since  $(y^n)_n$  weakly converges to 0 in  $L^2([0,T], H)$ , we see that  $([y^n])_n$  weakly converges to 0 in H. Indeed, for every  $\eta \in H$ , considering it as a constant function in  $L^2([0,T], H)$ , we have that

$$([y^n],\eta) = \left(\frac{1}{T}\int_0^T y^n(t) \, dt \, , \eta\right) = \frac{1}{T}\int_0^T (y^n(t) \, , \eta) \, dt \to 0 \, .$$

Moreover, by A5, the set  $\{w^n(t) : t \in [0,T], n \in \mathbb{N}\}$  is precompact in H. Hence, by Lemma 3.2(a), the sequence  $([w^n])_n$  is contained in a compact subset of H. Then, up to a subsequence,

$$\lim_{n} \left( [w^n], [y^n] \right) = 0 \,.$$

On the other hand, defining

$$\xi^{n}(t) = \int_{0}^{t} \check{w}^{n}(s) \, ds = (P\check{w}^{n})(t) \,,$$

we have that  $\xi^n(T) = \xi^n(0)$ , and recalling that  $\check{y}^n(t)$  and  $y^n(t)$  differ by a constant, integrating by parts we have

$$\int_0^T (\check{w}^n(t), \check{y}^n(t)) \, dt = -\int_0^T (\xi^n(t), \dot{y}^n(t)) \, dt \, .$$

We know that  $(\dot{y}^n)_n$  weakly converges to 0 in  $L^2([0,T], H)$ . Moreover, since  $\{w^n(t) : t \in [0,T], n \in \mathbb{N}\}$  is precompact in H, by Lemma 3.2(b), the sequence  $(\xi^n)_n$  is contained in a compact subset of C([0,T], H) and hence, up to a subsequence,

$$\lim_{n} \int_{0}^{T} (\xi^{n}(t), \dot{y}^{n}(t)) dt = 0,$$

thus proving (3.7). The Claim is thus proved.

Going back to (3.5), by (3.6) and (3.7), we get

$$\lim_{n} \int_{0}^{T} \left[ |\dot{z}^{n}(t) - \dot{z}^{*}(t)|^{2} - (\mathcal{A}(z^{n}(t) - z^{*}(t)), z^{n}(t) - z^{*}(t)) \right] dt = 0.$$

By A1, being  $\mathcal{A}$  semi-negative definite, we deduce that

$$\lim_{n} \int_{0}^{T} |\dot{z}^{n}(t) - \dot{z}^{*}(t)|^{2} dt = 0,$$

and

$$\lim_{n} \int_{0}^{T} (\mathcal{A}(z^{n}(t) - z^{*}(t)), z^{n}(t) - z^{*}(t)) dt = 0,$$

i.e.,

$$\lim_{n} \int_{0}^{T} \left[ (\mathcal{A}(\hat{z}^{n} - \hat{z}^{*}), \hat{z}^{n} - \hat{z}^{*}) + (\mathcal{A}(\check{z}^{n}(t) - \check{z}^{*}(t)), \check{z}^{n}(t) - \check{z}^{*}(t)) \right] dt = 0,$$

Hence,  $\dot{z}^n \to \dot{z}^*$  in  $L^2([0,T], H)$ , and, by A1, also  $\hat{z}^n \to \hat{z}^*$ . By Proposition 3.1,  $\dot{z}^n \to \dot{z}^*$  so that, being  $\tilde{z}^n = \hat{z}^n + \dot{z}^n$ , we have proved that  $(\tilde{z}^n)_n$  converges in  $H^1_T$ . This fact leads to the conclusion of the proof.

We now distinguish the two cases. If  $\mathcal{N}(\mathcal{A})$  has finite dimension N, then Theorem 2.2 applies, because Proposition 3.3 provides the  $(PS)_G$  condition for

$$G = \left\{ \sum_{k=1}^{N} m_k \tau_k a_k : m_k \in \mathbb{Z} \right\} \,,$$

which is a subgroup of  $H_T^1$ , and  $\varphi$  is *G*-invariant. We thus get N + 1 critical orbits of  $\varphi$ .

959

Assume now that  $\mathcal{N}(\mathcal{A})$  is infinite-dimensional. We first prove that  $\varphi$  has a minimum. To this aim, let  $(x^n)_n$  be a sequence in  $H^1_T$  such that  $\varphi(x^n) \to \iota := \inf \varphi(H^1_T)$ . By the Ekeland Principle, there is a sequence  $(y^n)_n$  such that

$$||x^n - y^n|| \to 0, \quad \varphi(y^n) \to \iota, \quad \nabla \varphi(y^n) \to 0.$$

Moreover, by (3.1), we can argue as in beginning of the proof of Proposition 3.3 and assume without loss of generality that

$$\bar{y}^n \in K := \left\{ y = \sum_{k=1}^{\infty} y_k a_k : y_k \in [0, \tau_k] \text{ for } k = 1, 2 \dots \right\} ,$$

for every *n*. The set *K* is compact, being isometric to the Hilbert cube  $\prod_{k=1}^{\infty} [0, \tau_k]$ in  $\ell^2$ , since  $\sum_{k=1}^{\infty} \tau_k^2 < +\infty$ . Using this and Proposition 3.3, there is a subsequence of  $(y^n)_n$  converging to some  $y^* \in H_T^1$ . Then,  $\varphi(y^*) = \iota$ , and  $\nabla \varphi(y^*) = 0$ . We have thus found a minimum point for the functional  $\varphi$ .

If  $y^*$  is not an isolated minimum point, then there are infinitely many minimum points near  $y^*$ . In this case, then, there are infinitely many geometrically distinct critical points of  $\varphi$ .

Otherwise, if  $y^*$  is an isolated minimum point, there is a constant r > 0 such that  $\varphi(u) > \min \varphi$ , for every  $u \in \overline{B}(y^*, r) \setminus \{y^*\}$ .

(We denote by  $B(y^*, r)$  the open ball centered at  $y^*$ , with radius r > 0, and by  $\overline{B}(y^*, r)$  its closure.) Let us prove that

$$\inf_{\partial B(y^*,r)} \varphi > \min \varphi \,.$$

By contradiction, assume that there is a sequence  $(\xi^n)_n$  in  $\partial B(y^*, r)$  such that  $\varphi(\xi^n) \to \min \varphi$ . Using the Ekeland Principle, it is possible to find a sequence  $(\eta^n)_n$  in  $H_T^1$  such that  $\varphi(\eta^n) \to \min \varphi$ ,  $\|\eta^n - \xi^n\| \to 0$  and  $\nabla \varphi(\eta^n) \to 0$ . By (3.1), we can assume without loss of generality that  $\bar{\eta}^n \in K$ , for every n. Then, by Proposition 3.3, there is a subsequence of  $(\eta^n)_n$  which converges to some y in  $H_T^1$ . Being  $\partial B(x, r)$  a closed set, we have that  $y \in \partial B(y^*, r)$ , and by continuity  $\varphi(y) = \min \varphi$ , a contradiction.

Choosing, e.g.,  $y^{**} = y^* + \tau_1 a_1$ , if r > 0 small enough we have that  $y^{**} \notin B(y^*, r)$ , and

$$\varphi(y^{**}) = \varphi(y^*) < \inf_{\partial B(y^*,r)} \varphi.$$

So, the Mountain Pass Theorem applies: setting

$$\Gamma = \{ \gamma \in C([0,1], H_T^1) : \gamma(0) = y^*, \gamma(1) = y^{**} \},\$$

and

$$c = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} \varphi(\gamma(t)) \,,$$

there is a sequence  $(u^n)_n$  in  $H^1_T$  such that

$$\lim_{n} \varphi(u^{n}) = c, \qquad \lim_{n} \nabla \varphi(u^{n}) = 0.$$

Moreover,  $\varphi(y^*) < c$ . Proceeding as in the first part of the proof, we can assume without loss of generality that  $\bar{u}^n \in K$ , and we can find a subsequence of  $(u^n)_n$  which

converges to some  $u^* \in H^1_T$ , so that  $\varphi(u^*) = c$  and  $\nabla \varphi(u^*) = 0$ . Since  $\varphi(y^*) < \varphi(u^*)$ , we have thus found two critical points,  $y^*$  and  $u^*$ , which are geometrically distinct.

### 4. Some examples and an open problem

Let  $(e_k)_k$  be an orthonormal basis in H, and assume that the periodicity condition (1.3) holds. Assume moreover (1.1), i.e., that e(t) has a zero mean. Defining, for every  $N \ge 1$ , the projection

$$P_N: H \to H$$
,  $x = \sum_{k=1}^{\infty} x_k e_k \mapsto \sum_{k=N+1}^{\infty} x_k e_k$ ,

we have that

$$\mathcal{N}(P_N) = \operatorname{span}\{e_1, \dots, e_N\}$$

Then, taking  $\mathcal{A} = -P_N$ , Theorem 2.1 applies to the system

$$\ddot{x} - P_N x + \nabla_x V(t, x) = e(t) \,,$$

and provides us with at least N + 1 geometrically distinct T-periodic solutions.

Notice that the number of T-periodic solutions increases indefinitely together with N. However, passing to the limit on N, the system becomes

$$\ddot{x} + \nabla_x V(t, x) = e(t) \,,$$

to which Theorem 2.1 still applies, but guarantees only two T-periodic solutions. It is an open problem to know if, in this last case, the existence of more than two T-periodic solutions can be proved.

As a first example of application, we consider the space  $H = \ell^2$  and the function

$$V(t,x) = -\sum_{k=1}^{+\infty} \frac{c_k}{\omega_k} \cos(\omega_k x_k) \cos(\omega_{k+1} x_{k+1}),$$

with  $c_k > 0$  and  $\omega_k > 0$ , for every  $k \ge 1$ . We have the cyclically coupled system

$$x_{k}'' + \left[\frac{c_{k-1}\omega_{k}}{\omega_{k-1}}\cos(\omega_{k-1}x_{k-1}) + c_{k}\cos(\omega_{k+1}x_{k+1})\right]\sin(\omega_{k}x_{k}) = e_{k}(t), \ k = 1, 2, \dots$$

where we have formally set  $c_0 = 0$  and  $\omega_0 = 1$ . Assuming that the sequences

$$(c_k)_k, \quad \left(\frac{1}{\omega_k}\right)_k, \quad \left(\frac{c_{k-1}\omega_k}{\omega_{k-1}}\right)_k$$

all belong to  $\ell^2$  (e.g., we could take  $c_k = 1/k$  and  $\omega_k = k$ ), we can apply Theorem 2.1, so that at least two *T*-periodic solutions exist.

Another example can be obtained if we now identify  $\ell^2$  with the space of sequences  $(\xi_k)_k$  where k ranges from  $-\infty$  to  $+\infty$ , i.e., with  $\ell^2(\mathbb{Z})$ . Defining

$$\mathcal{V}(t,x) = -\sum_{k=-\infty}^{+\infty} \frac{1}{\omega_k} \cos(\omega_k x_k) \Big( c'_k \cos(\omega_{k-1} x_{k-1}) + c''_k \cos(\omega_{k+1} x_{k+1}) \Big),$$

with  $c'_k, c''_k > 0$  and  $\omega_k > 0$  for every integer k, we have the system

$$x_k'' + [\alpha_k \cos(\omega_{k-1} x_{k-1}) + \beta_k \cos(\omega_{k+1} x_{k+1})] \sin(\omega_k x_k) = e_k(t), \quad k \in \mathbb{Z},$$

where

$$\alpha_k = \frac{c'_k \omega_{k-1} + c''_{k-1} \omega_k}{\omega_{k-1}}, \qquad \beta_k = \frac{c''_k \omega_{k+1} + c'_{k+1} \omega_k}{\omega_{k+1}}.$$

If we assume that all the sequences  $(c_k)_k$ ,  $(\omega_k^{-1})_k$ ,  $(\alpha_k)_k$ ,  $(\beta_k)_k$  belong to  $\ell^2(\mathbb{Z})$  (e.g., taking  $c'_k = c''_k = (|k|+1)^{-1}$  and  $\omega_k = |k|+1$ ), by Theorem 2.1 we conclude that at least two *T*-periodic solutions must exist.

Acknowledgement. The authors wish to thank Alberto Boscaggin and Maurizio Garrione for suggesting the two final examples, and the referee for valuable suggestions.

#### References

- A. Boscaggin, A. Fonda and M. Garrione, An infinite-dimensional version of the Poincaré-Birkhoff theorem on the Hilbert cube, Ann. Scuola Norm. Sup. Pisa Cl. Sci. 20 (2020), 751–770.
- H. Brezis, Opérateurs maximux monotones et semi-groupes de contractions dans les espaces de Hilbert, North-Holland Math. Studies 5, North-Holland, Amsterdam, 1973.
- [3] K.C. Chang, On the periodic nonlinearity and the multiplicity of solutions, Nonlinear Anal. 13 (1989), 527–537.
- [4] C.C. Conley and E.J. Zehnder, The Birkhoff-Lewis fixed point theorem and a conjecture of V.I. Arnold, Invent. Math. 73 (1983), 33-49.
- [5] E.N. Dancer, On the use of asymptotics in nonlinear boundary value problems, Ann. Mat. Pura Appl. 131 (1982), 167–185.
- [6] P.L. Felmer, Periodic solutions of spatially periodic Hamiltonian systems, J. Differential Equations 98 (1992), 143–168.
- [7] A. Fonda and J. Mawhin, Multiple periodic solutions of conservative systems with periodic nonlinearity, in: Differential equations and applications (Columbus, OH, 1988), 298–304, Ohio Univ. Press, Athens, OH, 1989.
- [8] A. Fonda and R. Toader, Periodic solutions of pendulum-like Hamiltonian systems in the plane, Adv. Nonlinear Stud. 12 (2012), 395–408.
- [9] A. Fonda and A.J. Ureña, A higher dimensional Poincaré-Birkhoff theorem for Hamiltonian flows, Ann. Inst. H. Poincaré Anal. Non Linéaire 34 (2017), 679–698.
- [10] G. Fournier, D. Lupo, M. Ramos and M. Willem, *Limit relative category and critical point theory*, in: Dynamics reported. Expositions in Dynamical Systems, vol. 3, 1–24, Springer, Berlin, 1994.
- [11] J. Franks, Generalizations of the Poincaré–Birkhoff theorem, Ann. Math. 128 (1988), 139–151.
- [12] G. Hamel, Über erzwungene Schwingungen bei endlichen Amplituden, Math. Ann. 86 (1922), 1–13.
- [13] F. Josellis, Lyusternik-Schnirelman theory for flows and periodic orbits for Hamiltonian systems on T<sup>n</sup> × ℝ<sup>n</sup>, Proc. London Math. Soc. (3) 68 (1994), 641–672.
- [14] J.Q. Liu, A generalized saddle point theorem, J. Differential Equations 82 (1989), 372–385.
- [15] J. Mawhin, Forced second order conservative systems with periodic nonlinearity, Analyse non linéaire (Perpignan, 1987), Ann. Inst. H. Poincaré Anal. Non Linéaire 6 (1989), suppl., 415– 434.
- [16] J. Mawhin and M. Willem, Multiple solutions of the periodic boundary value problem for some forced pendulum-type equations, J. Differential Equations 52 (1984), 264–287.
- [17] J. Mawhin and M. Willem, Variational methods and boundary value problems for vector second order differential equations and applications to the pendulum equation, in: Nonlinear Analysis and Optimization (Bologna, 1982), 181–192, Lecture Notes in Math. 1107, Springer, Berlin, 1984.

- [18] J. Mawhin and M. Willem, Critical point theory and Hamiltonian systems, Springer, Berlin, 1989.
- [19] P. Rabinowitz, On a class of functionals invariant under a  $\mathbb{Z}^n$  action. Trans. Amer. Math. Soc. **310** (1988), 303–311.
- [20] A. Szulkin, A relative category and applications to critical point theory for strongly indefinite functionals, Nonlinear Anal. 15 (1990), 725–739.
- [21] M. Willem, Oscillations forcées de l'équation du pendule, Publ. IRMA Besançon 3 (1981), v1-v3.

Manuscript received June 13 2019 revised October 21 2019

A. Fonda

Dipartimento di Matematica e Geoscienze, Università di Trieste, P.le Europa 1, I-34127 Trieste, Italy

*E-mail address*: a.fonda@units.it

J. Mawhin

Institut de Recherche en Mathématique et Physique, Université Catholique de Louvain, Chemin du Cyclotron 2, B-1348 Louvain-la-Neuve, Belgium

*E-mail address*: jean.mawhin@uclouvain.be

M. WILLEM

Institut de Recherche en Mathématique et Physique, Université Catholique de Louvain, Chemin du Cyclotron 2, B-1348 Louvain-la-Neuve, Belgium

*E-mail address*: michel.willem@uclouvain.be