

MÖBIUS ORTHOGONALITY IN DENSITY FOR ZERO ENTROPY DYNAMICAL SYSTEMS

ALEXANDER GOMILKO, MARIUSZ LEMAŃCZYK, AND THIERRY DE LA RUE

ABSTRACT. It is proved that whenever a zero entropy dynamical system (X, T) has only countably many ergodic measures and μ stands for the arithmetic Möbius function then there exists $A = A(X, T) \subset \mathbb{N}$ of logarithmic density one such that for each $f \in C(X)$,

$$\lim_{A \ni N \rightarrow \infty} \frac{1}{N} \sum_{n \leq N} f(T^n x) \mu(n) = 0$$

uniformly in $x \in X$, in particular, the density version of Möbius orthogonality holds.

1. INTRODUCTION

Following P. Sarnak [14], we say that a topological system (X, T) is *Möbius orthogonal* if

$$(1.1) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{n \leq N} f(T^n x) \mu(n) = 0$$

for all $f \in C(X)$ and $x \in X$ (here μ stands for the classical arithmetic Möbius function). By the standard trick of summation by parts (which we recall below for the reader's convenience), we obtain that the Möbius orthogonality of (X, T) implies the *logarithmic Möbius orthogonality* of (X, T) :

$$(1.2) \quad \lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N} \frac{1}{n} f(T^n x) \mu(n) = 0$$

for all $f \in C(X)$ and $x \in X$. The celebrated Sarnak's conjecture [14] claims that all zero entropy systems are Möbius orthogonal, but this statement has been established only for some selected classes (we refer the reader to the bibliography in survey [9] to see for which classes). In some contrast to this, a considerable progress has been made recently in our understanding of the logarithmic Sarnak's conjecture: Frantzikinakis and Host [10] proved that each zero entropy system whose set of ergodic measures is countable is logarithmic Möbius orthogonal. Earlier, Tao [17] proved that the logarithmic Sarnak's conjecture is equivalent to the logarithmic version of the classical Chowla conjecture (from 1965) on auto-correlations of the Möbius function.¹

¹Sarnak's conjecture itself was motivated by the fact that the Chowla conjecture implies Sarnak's conjecture [2], [14], [15]. See also [16], [19], [20], where special cases of the validity of the logarithmic Chowla conjecture have been proved.

While it looked rather odd to expect that we can say anything interesting about Cesàro type averaging knowing the convergence of logarithmic averages, it has been proved in [12] that the logarithmic Chowla conjecture implies the validity of the Chowla conjecture along a subsequence. In fact, the result was a consequence of some general mechanism when a certain sequence (in a locally convex space) of logarithmic averages² converges to an extremal point. This approach seems to fail if (what perhaps is natural), we would like to prove that the logarithmic Möbius orthogonality of a fixed system implies its Möbius orthogonality along a subsequence. However, commenting on [12], Tao [18], using a different method (second moment type argument), was able to prove a stronger result, namely, he proved that if the logarithmic Chowla conjecture holds then the Chowla conjecture holds along a subsequence of full logarithmic density; in particular, of upper density 1.

The aim of this note is to show how to adapt Tao's argument (this is done in Theorem 2.1 below) to be able to apply it to systems satisfying some (seemingly) stronger condition than the logarithmic Möbius orthogonality and which allows one to deduce Möbius orthogonality in full logarithmic density. In order to formulate such a result we first recall the *strong MOMO* notion introduced in [3]. Namely, a system (X, T) satisfies this property if for all increasing sequences $(b_k) \subset \mathbb{N}$ with $b_{k+1} - b_k \rightarrow \infty$, all $(x_k) \subset X$ and $f \in C(X)$, we have

$$(1.3) \quad \frac{1}{b_{K+1}} \sum_{k \leq K} \left| \sum_{b_k \leq n < b_{k+1}} f(T^{n-b_k} x_k) \mu(n) \right| \rightarrow 0,$$

while if

$$(1.4) \quad \frac{1}{\log b_{K+1}} \sum_{k \leq K} \left| \sum_{b_k \leq n < b_{k+1}} \frac{1}{n} f(T^{n-b_k} x_k) \mu(n) \right| \rightarrow 0,$$

then we say that (X, T) satisfies the *logarithmic strong MOMO* property. It has been proved in [3] that Sarnak's conjecture is equivalent to the fact that all zero entropy systems enjoy the strong MOMO property.

Note that (1.3) is equivalent to

$$(1.5) \quad \lim_{K \rightarrow \infty} \frac{1}{b_{K+1}} \sum_{k \leq K} \left\| \sum_{b_k \leq n < b_{k+1}} \mu(n) f \circ T^n \right\|_{C(X)} = 0,$$

and (1.4) is equivalent to

$$(1.6) \quad \lim_{K \rightarrow \infty} \frac{1}{\log b_{K+1}} \sum_{k \leq K} \left\| \sum_{b_k \leq n < b_{k+1}} \frac{\mu(n)}{n} f \circ T^n \right\|_{C(X)} = 0,$$

²The Chowla conjecture can be reformulated using the language of quasi-generic points for invariant measures in a certain shift space; it is then equivalent to the fact that the empiric measures determined by μ converge to a certain natural measure which is ergodic, hence to an extremal point, see e.g. the survey [9]. Similarly, we deal with the logarithmic Chowla conjecture.

for all increasing sequences $(b_k) \subset \mathbb{N}$ with $b_{k+1} - b_k \rightarrow \infty$ and $f \in C(X)$. From (1.5) we obtain

$$\lim_{k \rightarrow \infty} \frac{1}{b_{k+1}} \left\| \sum_{n < b_{k+1}} \mu(n) f \circ T^n \right\|_{C(X)} = 0,$$

for $b_{k+1} - b_k \rightarrow \infty$, $k \rightarrow \infty$, and then it is easy to see that it holds for $b_k := k$, such that the uniform convergence in Möbius orthogonality (1.1) holds. By analogous way we obtain that logarithmic strong MOMO property implies the uniform convergence in logarithmic Möbius orthogonality (1.2).

Here is our main result:

Theorem 1.1. *Assume that a topological system (X, T) satisfies the logarithmic strong MOMO property. Then there exists $A = A(X, T) \subset \mathbb{N}$ with full logarithmic density: $\delta(A) = 1$, such that for each $f \in C(X)$,*

$$(1.7) \quad \lim_{A \ni N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n \leq N} \mu(n) f \circ T^n \right\|_{C(X)} = 0.$$

In particular, Möbius orthogonality holds along a subsequence (of N) of full logarithmic density.

One of the main results in [10] states that, if a system (X, T) has zero entropy and if its set of ergodic measures is countable, then the system is logarithmic Möbius orthogonal. We will show that such systems satisfy the strong logarithmic MOMO property, hence obtaining the following:³

Corollary 1.2. *Let (X, T) be a zero entropy ergodic dynamical system such that the set $M^e(X, T)$ of ergodic T -invariant measures is countable. Then, there exists $A = A(X, T) \subset \mathbb{N}$ with full logarithmic density along which Möbius orthogonality holds uniformly in $x \in X$.*

In particular, the above holds for all zero entropy uniquely ergodic systems. Corollary 1.2 is slightly surprising even for horocycle flows (in the cocompact case), where we know that Möbius orthogonality holds [5] but it is open (see questions in [9], [13]) whether Möbius orthogonality holds in its uniform form. By Corollary 1.2, we have that a uniform version holds along a subsequence of logarithmic density 1 (let alone the upper density of this subsequence is 1). A use of [11] shows that Corollary 1.2 remains valid when μ is replaced by any multiplicative function which is strongly aperiodic.

The rest of the note is devoted to give some illustrations how Theorem 2.1 (which is an adaptation of Tao's result) can be applied in other situations. For example, we will show how in the main result in [12] we can achieve full logarithmic density. Besides, we note that in the classical Davenport-Erdős theorem on the existence of the logarithmic density [7] of sets of multiples, the upper asymptotic density is achieved along a set of full logarithmic density. Finally, we note in passing the logical

³We were informed by N. Frantzikinakis during the workshop "Sarnak's Conjecture" at the American Institute of Mathematics in mid-December 2018 that, independently of us, he can prove Corollary 1.2 by modifying some arguments in [10].

implication: logarithmic Chowla conjecture of order 2 \Rightarrow PNT along a subsequence of logarithmic density 1.

A few words on basic concepts and notation: Given a subset $C \subset \mathbb{N}$, we denote by $\delta(C)$ its logarithmic density: $\delta(C) := \lim_{N \rightarrow \infty} (1/\log N) \sum_{n \leq N, n \in C} \frac{1}{n}$, assuming that the limit exists. It is classical that $\underline{d}(C) \leq \delta(C) \leq \bar{d}(C)$, where $\bar{d}(C) := \limsup_{N \rightarrow \infty} \frac{1}{N} |[1, N] \cap C|$ stands for the upper asymptotic density, and similarly the lower (asymptotic) density $\underline{d}(C)$ is defined as the lim inf. In fact, these inequalities are direct consequences of the classical relationship between Cesàro averages and logarithmic averages: given a sequence (a_n) and setting $s_n := \sum_{j \leq n} a_j$, $s_0 := 0$, we have:

$$(1.8) \quad \sum_{1 \leq n \leq N} \frac{a_n}{n} = \sum_{1 \leq n \leq N} \frac{1}{n} (s_n - s_{n-1}) = \sum_{1 \leq n \leq N-1} s_n \left(\frac{1}{n} - \frac{1}{n+1} \right) + \frac{s_N}{N} = \sum_{n \leq N-1} \frac{s_n}{n} \frac{1}{n+1} + \frac{s_N}{N}$$

which basically says that the logarithmic averages of (a_n) are the logarithmic averages of Cesàro averages (sometimes, we only use the fact that the logarithmic averages are convex combinations of Cesàro averages).

In what follows when we speak about subsequences of natural numbers, we always mean increasing sequences of natural numbers (so that subsequences are the same as infinite subsets). In Corollary 1.2, we find a subsequence of full logarithmic density which depends however on the system (X, T) . The methods used in this note do not seem to give one universal subsequence along which Sarnak's conjecture (i.e. Möbius orthogonality for zero entropy systems) holds. We could get such a universal sequence (see Proposition 1.3 below) if we were able to prove Sarnak's conjecture along a full logarithmic density sequence for **each** zero entropy system, that is, by [17], if the logarithmic Chowla conjecture holds. More precisely:

Proposition 1.3. *Assume that for each zero entropy dynamical system (X, T) there exists a subsequence $(N_k(X, T))_k$ of natural numbers with $\delta(\{N_k(X, T) : k \geq 1\}) = 1$ such that*

$$(1.9) \quad \lim_{k \rightarrow \infty} \frac{1}{N_k(X, T)} \sum_{n \leq N_k(X, T)} f(T^n x) \mu(n) = 0$$

for all $f \in C(X)$ and $x \in X$. Then there exists a subsequence (N_k) of natural numbers, $\delta(\{N_k : k \geq 1\}) = 1$ such that for each zero entropy dynamical system (X, T) , (1.9) holds along (N_k) .

To see the proof of Proposition 1.3, we have:

a) By assumption and the classical Lemma 2.5, we obtain that for each zero entropy dynamical system (X, T) , we have $\lim_{N \rightarrow \infty} (1/\log N) \sum_{n \leq N} \frac{1}{n} f(T^n x) \mu(n) = 0$ for each $f \in C(X)$ and $x \in X$.

(b) By (a) and Tao's result ("logarithmic Sarnak implies logarithmic Chowla") [17], in the space $M(X_\mu)$ of measures on X_μ , we obtain that $(1/\log N) \sum_{n \leq N} (1/n) \delta_{S^n \mu} \rightarrow \hat{\nu}_\mathcal{S}$, where we consider the Möbius subshift (X_μ, S) and $\hat{\nu}_\mathcal{S}$

stands for the relatively independent extension of the Mirsky measure $\nu_{\mathcal{S}}$ of the square-free system (X_{μ^2}, S) .⁴

(c) By (b) and Theorem 5.1, we obtain that there exists a subsequence (N_k) with $\delta(\{N_k : k \geq 1\}) = 1$ such that $(1/N_k) \sum_{n \leq N_k} \delta_{S^n \mu} \rightarrow \hat{\nu}_{\mathcal{S}}$.

(d) By (c) and the proof of the implication “Chowla implies Sarnak” in [2], it follows that for each zero entropy (X, T) , we have $\lim_{k \rightarrow \infty} (1/N_k) \sum_{n \leq N_k} f(T^n x) \mu(n) = 0$ for all $f \in C(X)$ and $x \in X$, so Proposition 1.3 follows.

2. FUNCTIONAL FORMULATION OF TAO’S RESULT

Our aim in this section is to prove a slight extension of Tao’s result from [18]:

Theorem 2.1. *Let $(\mathcal{B}_j, \|\cdot\|_j)$, $j = 1, 2$, be normed vector spaces and assume that \mathcal{B}_1 is separable. Let $(S_k)_{k \geq 1}$ be a sequence of linear bounded operators from \mathcal{B}_1 to \mathcal{B}_2 , such that for some M we have, for each $f \in \mathcal{B}_1$ and each $k \geq 1$,*

$$(2.1) \quad \|S_k f\|_2 \leq M \|f\|_1. \quad^5$$

Let ϕ be a continuous, positive, strictly increasing and convex function on $[0, \infty)$ with $\phi(0) = 0$. Suppose that there exists a subsequence $(N_s) \subset \mathbb{N}$ such that for any $f \in \mathcal{B}_1$, setting

$$(2.2) \quad R_f(H) := \limsup_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{1 \leq n \leq N_s} \frac{1}{n} \phi \left(\left\| \frac{1}{H} \sum_{1 \leq h \leq H} S_{n+h} f \right\|_2 \right),$$

we have

$$(2.3) \quad \lim_{H \rightarrow \infty} R_f(H) = 0.$$

Then there exists a set \mathcal{N} of natural numbers with the property

$$(2.4) \quad \lim_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{\mathcal{N} \ni N \leq N_s} \frac{1}{N} = 1,$$

such that, for any $f \in \mathcal{B}_1$, we have

$$(2.5) \quad \lim_{N \rightarrow \infty, N \in \mathcal{N}} \left\| \frac{1}{N} \sum_{1 \leq n \leq N} S_n f \right\|_2 = 0.$$

Moreover, if $(N_s) = \mathbb{N}$, then $\delta(\mathcal{N}) = 1$ and

$$(2.6) \quad \lim_{N \rightarrow \infty} \left\| \frac{1}{\log N} \sum_{1 \leq n \leq N} \frac{S_n f}{n} \right\|_2 = 0.$$

The proof of the above theorem requires a few lemmas.

⁴The measure-theoretic investigations of the square-free system $(X_{\mu^2}, \nu_{\mathcal{S}}, S)$ have been originated by Sarnak [14] and Cellarosi and Sinai [6]: the Mirsky measure is ergodic and so is its relatively independent extension.

⁵Assuming that \mathcal{B}_1 is Banach and using the Uniform Boundedness Principle, we only need to assume that in (2.1) we have $\sup_{k \in \mathbb{N}} \|S_k f\|_2 < +\infty$.

Lemma 2.2. *Let $G : \mathbb{N} \rightarrow \mathbb{R}_+$. Suppose that for some $\gamma \in (0, 1)$ and for some subsequence $(N_s) \subset \mathbb{N}$, we have*

$$(2.7) \quad \limsup_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{1 \leq n \leq N_s} \frac{G(n)}{n} \leq \gamma.$$

Then, for the set $M \subset \mathbb{N}$ given by $M := \{n : G(n) < \sqrt{\gamma}\}$, we have

$$(2.8) \quad \liminf_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{n \leq N_s, n \in M} \frac{1}{n} \geq 1 - \sqrt{\gamma}.$$

Proof. Let $Q = \mathbb{N} \setminus M$, so that $Q = \{n : G(n) \geq \sqrt{\gamma}\}$. By Markov's inequality, we obtain

$$\begin{aligned} \frac{1}{\log N_s} \sum_{1 \leq n \leq N_s} \frac{G(n)}{n} &\geq \frac{\sqrt{\gamma}}{\log N_s} \sum_{n \in Q, n \leq N_s} \frac{1}{n} = \\ &= \frac{\sqrt{\gamma}}{\log N_s} \sum_{n \leq N_s} \frac{1}{n} - \frac{\sqrt{\gamma}}{\log N_s} \sum_{n \in M, n \leq N_s} \frac{1}{n}, \end{aligned}$$

hence

$$\frac{1}{\log N_s} \sum_{n \in M, n \leq N_s} \frac{1}{n} \geq \frac{1}{\log N_s} \sum_{n \leq N_s} \frac{1}{n} - \frac{1}{\sqrt{\gamma} \log N_s} \sum_{1 \leq n \leq N_s} \frac{G(n)}{n},$$

and (2.8) holds in view of (2.7). \square

Lemma 2.3. *Assume that $F : \mathbb{N} \rightarrow \mathbb{R}_+$ is bounded and satisfies*

$$(2.9) \quad \limsup_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{1 \leq n \leq N_s} \frac{F(n)}{n} \leq \gamma, \quad \gamma \in (0, 1),$$

for a subsequence (N_s) . Then, for the set $T \subset \mathbb{N}$ given by

$$(2.10) \quad T := \left\{ N : \frac{1}{N} \sum_{1 \leq n \leq N} F(n) \leq \sqrt{\gamma} \right\},$$

we have

$$(2.11) \quad \liminf_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{n \in T, n \leq N_s} \frac{1}{n} \geq 1 - \sqrt{\gamma}.$$

Proof. Set $G(n) := \frac{1}{n} \sum_{1 \leq m \leq n} F(m)$. By (1.8), we have

$$\sum_{1 \leq n \leq N-1} \frac{G(n)}{n+1} = \sum_{1 \leq n \leq N} \frac{F(n)}{n} - \frac{1}{N} \sum_{1 \leq n \leq N} F(n),$$

so that, by (2.9), we have

$$\limsup_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{1 \leq n \leq N_s} \frac{G(n)}{n} = \limsup_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{1 \leq n \leq N_s-1} \frac{G(n)}{n+1} \leq \gamma.$$

Then, by Lemma 2.2, we obtain (2.11) for the set T defined by (2.10). \square

Lemma 2.4 (see Tao [18]). Assume that $M_k \subset \mathbb{N}$, $k \in \mathbb{N}$, and that there exists an increasing sequence $(N_s) \subset \mathbb{N}$ such that, for each k ,

$$\lim_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{n \in M_k, n \leq N_s} \frac{1}{n} = 1, \quad k \in \mathbb{N}.$$

Then there exists a subset $M \subset \mathbb{N}$ such that

$$(2.12) \quad \lim_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{n \in M, n \leq N_s} \frac{1}{n} = 1,$$

and such that, for any $k \in \mathbb{N}$, there exists s_k with

$$(2.13) \quad M \cap \{n : n \geq N_{s_k}\} \subset M_k.$$

Proof. Replacing, if necessary, each M_k by $M_k \cap \bigcap_{k' < k} M_{k'}$, we may assume without loss of generality that $M_{k+1} \subset M_k$, $k \in \mathbb{N}$.

Let us choose an increasing sequence (s_k) such that, for each k ,

$$s \geq s_k \Rightarrow \frac{1}{\log N_s} \sum_{n \in M_k, n \leq N_s} \frac{1}{n} \geq 1 - \frac{1}{k}.$$

We set $M := M_1 \cap [1, N_{s_1}] \cup \bigcup_{k=2}^{\infty} (M_k \cap (N_{s_{k-1}}, N_{s_k}])$ and verify that M satisfies the desired properties (2.12) and (2.13). \square

The following is classical.

Lemma 2.5. Let $(a_n)_{n \geq 1}$ be a bounded sequence in a normed vector space $(\mathcal{B}, \|\cdot\|)$. Suppose that there exists a subsequence $\mathcal{N} \subset \mathbb{N}$, $\delta(\mathcal{N}) = 1$, such that

$$(2.14) \quad \lim_{N \rightarrow \infty, N \in \mathcal{N}} \left\| \frac{1}{N} \sum_{1 \leq n \leq N} a_n \right\| = 0.$$

Then

$$(2.15) \quad \lim_{N \rightarrow \infty} \left\| \frac{1}{\log N} \sum_{1 \leq n \leq N} \frac{a_n}{n} \right\| = 0.$$

Proof. Let $\mathcal{M} = \mathbb{N} \setminus \mathcal{N}$, so that

$$(2.16) \quad \lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N, n \in \mathcal{M}} \frac{1}{n} = 0.$$

By (1.8), we have

$$\sum_{1 \leq n \leq N} \frac{a_n}{n} = \mathcal{E}_N + \sum_{1 \leq n \leq N-1} \frac{\mathcal{E}_n}{n+1}, \quad \text{where } \mathcal{E}_n := \frac{1}{n} \sum_{1 \leq m \leq n} a_m.$$

Setting $C := \sup_n \|\mathcal{E}_n\|$, we obtain

$$\begin{aligned} \left\| \frac{1}{\log N} \sum_{1 \leq n \leq N} \frac{a_n}{n} \right\| &\leq \frac{C}{\log N} + \left\| \frac{1}{\log N} \sum_{n \leq N, n \in \mathcal{M}} \frac{\mathcal{E}_n}{n+1} \right\| + \left\| \frac{1}{\log N} \sum_{n \leq N, n \in \mathcal{N}} \frac{\mathcal{E}_n}{n+1} \right\| \leq \\ &\frac{C}{\log N} + \frac{C}{\log N} \sum_{n \leq N, n \in \mathcal{M}} \frac{1}{n} + \left\| \frac{1}{\log N} \sum_{n \leq N, n \in \mathcal{N}} \frac{\mathcal{E}_n}{n+1} \right\|, \end{aligned}$$

and then assertion (2.15) follows from (2.16) and (2.14). \square

Proof of Theorem 2.1. Let $f \in \mathcal{B}_1$. By (2.2), (2.3) and Lemma 2.3, we obtain that for any fixed large $H \in \mathbb{N}$, there exists a set $\mathcal{N}_{f,H}$ with the property

$$\liminf_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{\mathcal{N}_{f,H} \ni N \leq N_s} \frac{1}{N} \geq 1 - \sqrt{R_f(H)},$$

and such that, for $N \in \mathcal{N}_{f,H}$, we have

$$\frac{1}{N} \sum_{1 \leq n \leq N} \phi \left(\left\| \frac{1}{H} \sum_{1 \leq h \leq H} S_{n+hf} \right\|_2 \right) \leq \sqrt{R_f(H)}.$$

By deleting at most finitely many elements, we may assume that $\mathcal{N}_{f,H}$ consists only of elements of size at least H^2 . For any H_0 , if we set $\mathcal{N}_{f, \geq H_0} := \bigcup_{H \geq H_0} \mathcal{N}_{f,H}$, then $\mathcal{N}_{f, \geq H_0}$ satisfies $\lim_{s \rightarrow \infty} (1/\log N_s) \sum_{\mathcal{N}_{f, \geq H_0} \ni N \leq N_s} \frac{1}{N} = 1$. By Lemma 2.4, we can find a set \mathcal{N}_f of natural numbers with

$$\lim_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{\mathcal{N}_f \ni N \leq N_s} \frac{1}{N} = 1,$$

and such that, for every H_0 , every sufficiently large element of \mathcal{N}_f lies in $\mathcal{N}_{f, \geq H_0}$. Thus, for every sufficiently large $N \in \mathcal{N}_f$, one has

$$\frac{1}{N} \sum_{1 \leq n \leq N} \phi \left(\left\| \frac{1}{H} \sum_{1 \leq h \leq H} S_{n+hf} \right\|_2 \right) \leq \sqrt{R_f(H)},$$

for some $H \geq H_0$ with $N \geq H^2$. By the monotonicity of ϕ and Jensen's inequality, this implies that

$$\begin{aligned} \phi \left(\frac{1}{NH} \left\| \sum_{1 \leq n \leq N} \sum_{1 \leq h \leq H} S_{n+hf} \right\|_2 \right) &\leq \phi \left(\frac{1}{N} \sum_{1 \leq n \leq N} \left\| \frac{1}{H} \sum_{1 \leq h \leq H} S_{n+hf} \right\|_2 \right) \\ &\leq \frac{1}{N} \sum_{1 \leq n \leq N} \phi \left(\left\| \frac{1}{H} \sum_{1 \leq h \leq H} S_{n+hf} \right\|_2 \right) \leq \sqrt{R_f(H)}, \end{aligned}$$

so that, setting $\psi(s) := \phi^{-1}(\sqrt{s})$, $s > 0$, we get

$$(2.17) \quad \frac{1}{NH} \left\| \sum_{1 \leq n \leq N} \sum_{1 \leq h \leq H} S_{n+hf} \right\|_2 \leq \psi(R_f(H)) \rightarrow 0, \quad H \rightarrow \infty.$$

Next, by computation, we get

$$\begin{aligned} \sum_{1 \leq n \leq N} \sum_{1 \leq h \leq H} S_{n+hf} &= \sum_{1 \leq h \leq H} \sum_{1 \leq n \leq N} S_{n+hf} = \sum_{1 \leq h \leq H} \sum_{m=h+1}^{h+N} S_m f = \\ &= H \sum_{m=1}^N S_m f + \sum_{1 \leq h \leq H} \sum_{m=N+1}^{N+h} S_m f - \sum_{1 \leq h \leq H} \sum_{1 \leq m \leq h} S_m f, \end{aligned}$$

which gives

$$\begin{aligned} \frac{1}{N} \sum_{m=1}^N S_m f &= \frac{1}{HN} \sum_{1 \leq n \leq N} \sum_{1 \leq h \leq H} S_{n+h} \\ &\quad - \frac{1}{HN} \sum_{1 \leq h \leq H} \sum_{m=N+1}^{N+h} S_m f + \frac{1}{HN} \sum_{1 \leq h \leq H} \sum_{1 \leq m \leq h} S_m f. \end{aligned}$$

Now, let us fix H_0 . For any sufficiently large $N \in \mathcal{N}_f$, there exists $H \geq H_0$, with $H^2 \leq N$, such that $N \in \mathcal{N}_{f,H}$. The triangular inequality yields

$$\begin{aligned} \frac{1}{N} \left\| \sum_{m=1}^N S_m f \right\|_2 &\leq \frac{1}{HN} \left\| \sum_{1 \leq n \leq N} \sum_{1 \leq h \leq H} S_{n+h} f \right\|_2 \\ &\quad + \frac{1}{HN} \left\| \sum_{1 \leq h \leq H} \sum_{m=N+1}^{N+h} S_m f \right\|_2 + \frac{1}{HN} \left\| \sum_{1 \leq h \leq H} \sum_{1 \leq m \leq h} S_m f \right\|_2. \end{aligned}$$

Using (2.17), we can upper bound the first term in the right-hand side by $\psi(R_f(H))$. The sum of the other two terms can be upper bounded by

$$\frac{M\|f\|_1}{HN} \left(\sum_{1 \leq h \leq H} \sum_{m=N+1}^{N+h} 1 + \sum_{1 \leq h \leq H} \sum_{1 \leq m \leq h} 1 \right) \leq \frac{2HM\|f\|_1}{N} \leq \frac{2M\|f\|_1}{H}.$$

We thus get that

$$\frac{1}{N} \left\| \sum_{m=1}^N S_m f \right\|_2 \leq \sup_{H \geq H_0} \psi(R_f(H)) + \frac{2M\|f\|_1}{H_0}.$$

By letting H_0 go to infinity, we conclude that $\lim_{N \rightarrow \infty, N \in \mathcal{N}_f} \left\| (1/N) \sum_{m=1}^N S_m f \right\|_2 = 0$.

Let now $(f_k)_{k \geq 1}$ be a dense set in \mathcal{B}_1 . Then, by Lemma 2.4, we get a set \mathcal{N} of natural numbers with the property $\lim_{s \rightarrow \infty} (1/\log N_s) \sum_{N \in \mathcal{N}, N \leq N_s} \frac{1}{N} = 1$, such that for any $k \in \mathbb{N}$,

$$\lim_{N \rightarrow \infty, N \in \mathcal{N}} \frac{1}{N} \left\| \sum_{m=1}^N S_m f_k \right\|_2 = 0.$$

Take $g \in \mathcal{B}_1$. Then for any $\epsilon > 0$ there exists f_k such that $\|g - f_k\|_1 \leq \epsilon/M$, and then

$$\frac{1}{N} \left\| \sum_{m=1}^N S_m g \right\|_2 \leq \frac{1}{N} \left\| \sum_{m=1}^N S_m f_k \right\|_2 + \epsilon,$$

which, by the above, yields

$$\lim_{N \rightarrow \infty, N \in \mathcal{N}} \frac{1}{N} \left\| \sum_{m=1}^N S_m g \right\|_2 = 0.$$

So, statement (2.5) is proved. Assertion (2.6) follows from (2.5) by Lemma 2.5. \square

Note that in the sequel we will use Theorem 2.1 mainly for $\phi(s) = s$ (or sometimes for $\phi(s) = s^2$).

3. PROOF OF THEOREM 1.1

To prove Theorem 1.1 we will show a stronger result:

Theorem 3.1. *Let $(\mathcal{B}_j, \|\cdot\|_j)$, $j = 1, 2$, be normed vector spaces and assume that \mathcal{B}_1 is separable. Let $(S_k)_{k \geq 1}$ be a sequence of linear bounded operators from \mathcal{B}_1 to \mathcal{B}_2 , such that for some $M > 0$ we have, for each $f \in \mathcal{B}_1$ and each $k \geq 1$,*

$$\|S_k f\|_2 \leq M \|f\|_1.$$

Assume that $(S_k)_{k \geq 1}$ satisfies the property: for all increasing sequences $(b_k) \subset \mathbb{N}$ with $b_{k+1} - b_k \rightarrow \infty$ and $f \in \mathcal{B}_1$, we have

$$\lim_{K \rightarrow \infty} \frac{1}{\log b_{K+1}} \sum_{k \leq K} \left\| \sum_{b_k \leq n < b_{k+1}} \frac{1}{n} S_n f \right\|_2 = 0,$$

Then there exists $A \subset \mathbb{N}$ with full logarithmic density: $\delta(A) = 1$, such that for each $f \in \mathcal{B}_1$,

$$\lim_{A \ni N \rightarrow \infty} \left\| \frac{1}{N} \sum_{n \leq N} S_n f \right\|_2 = 0.$$

The proof of Theorem 3.1 will use the following intermediate result whose proof depends on some ideas which first appeared in [1].

Proposition 3.2. *Let $(\mathcal{B}_j, \|\cdot\|_j)$, $j = 1, 2$ be normed vector spaces. Let $(S_n)_{n \geq 1}$ be a sequence of linear bounded operators from \mathcal{B}_1 to \mathcal{B}_2 , such that for any $(b_k) \subset \mathbb{N}$ with $0 < b_{k+1} - b_k \rightarrow \infty$ as $k \rightarrow \infty$, and any $f \in \mathcal{B}_1$ we have*

$$(3.1) \quad \frac{1}{\log b_{K+1}} \sum_{k \leq K} \left\| \sum_{b_k \leq n < b_{k+1}} \frac{1}{n} S_n f \right\|_2 \xrightarrow{K \rightarrow \infty} 0.$$

Then

$$(3.2) \quad \overline{\lim}_{H \rightarrow \infty} \overline{\lim}_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N} \frac{1}{n} \left\| \frac{1}{H} \sum_{h \leq H} S_{n+h} f \right\|_2 = 0.$$

We will prove this result by contraposition, and we will need the following lemma.

Lemma 3.3 (Diagonalization Lemma). *Consider a family of sequences $(g_{n,m}) \subset \mathbb{R}_+$, $m > n$, $m, n \geq 1$. Suppose that for some families $(b_{k,\ell})_{k,\ell \geq 1} \subset \mathbb{N}$ with*

$$(3.3) \quad 0 < b_{k,\ell} < b_{k+1,\ell}, \quad \overline{\lim}_{\ell \rightarrow \infty} \underline{\lim}_{k \rightarrow \infty} (b_{k+1,\ell} - b_{k,\ell}) = \infty,$$

and some $\gamma > 0$, we have

$$(3.4) \quad \overline{\lim}_{K \rightarrow \infty} \frac{1}{\log b_{K+1,\ell}} \sum_{k=1}^K g_{b_{k,\ell}, b_{k+1,\ell}} \geq \gamma, \quad \text{for all } \ell \in \mathbb{N}.$$

Then there exists a sequence $(b_k)_{k \geq 1} \subset \mathbb{N}$ such that

$$(3.5) \quad 0 < b_k < b_{k+1}, \quad b_{k+1} - b_k \xrightarrow{k \rightarrow \infty} \infty,$$

and

$$(3.6) \quad \overline{\lim}_{K \rightarrow \infty} \frac{1}{\log b_{K+1}} \sum_{k=1}^K g_{b_k, b_{k+1}} \geq \gamma/2.$$

Proof. Note that by (3.3), we have $b_{k+1,1} - b_{k,1} \geq 1$ for each $k \geq 1$, and that without loss of generality we may assume that for each $\ell \geq 1$, we have $\underline{\lim}_{k \rightarrow \infty} (b_{k+1,\ell} - b_{k,\ell}) \geq \ell$. By (3.4) for $\ell = 1$, we can choose the value K_1 so that

$$\frac{1}{\log b_{K_1+1,1}} \sum_{k=1}^{K_1} g_{b_{k,1}, b_{k+1,1}} \geq \gamma/2$$

and take then, for $k = 1, \dots, K_1 + 1$, $b_k := b_{k,1}$, to obtain

$$\frac{1}{\log b_{K_1+1}} \sum_{k=1}^{K_1} g_{b_k, b_{k+1}} \geq \gamma/2.$$

We continue the above process by induction. Suppose that for some $\ell \geq 1$ we already have sequences $0 = K_0 < K_1 < \dots < K_\ell$, $1 \leq b_1 < b_2 < \dots < b_{K_\ell+1}$, satisfying, for each $s = 1, \dots, \ell$,

$$(3.7) \quad b_{k+1} - b_k \geq s, \quad k = K_{s-1} + 1, \dots, K_s,$$

and

$$(3.8) \quad \frac{1}{\log b_{K_s+1}} \sum_{k=1}^{K_s} g_{b_k, b_{k+1}} \geq \gamma/2.$$

Then, at step $\ell + 1$, we first choose $N_{\ell+1}$ large enough so that $b_{K_\ell+2+N_{\ell+1}, \ell+1} \geq b_{K_\ell+1} + \ell + 1$, and, for each $k \geq K_\ell + 2 + N_{\ell+1}$, $b_{k+1, \ell+1} - b_{k, \ell+1} \geq \ell + 1$. Then, by (3.4), we can choose $K_{\ell+1} > K_\ell + 2$ large enough so that

$$\frac{1}{\log b_{K_{\ell+1}+1+N_{\ell+1}, \ell+1}} \sum_{k=K_\ell+2+N_{\ell+1}}^{K_{\ell+1}+N_{\ell+1}} g_{b_{k, \ell+1}, b_{k+1, \ell+1}} \geq \gamma/2,$$

and we take for $k = K_\ell + 2, \dots, K_{\ell+1} + 1$:

$$b_k := b_{k+N_{\ell+1}, \ell+1}.$$

Then assertions (3.7) and (3.8) are valid up to $s = \ell + 1$, and this allows us to construct inductively the sequences $(b_k)_{k \geq 1}$ with the required properties (3.5), (3.6). \square

Proof of Proposition 3.2. Suppose that for some $f \in \mathcal{B}_1$, (3.2) does not hold. Then, there exist $\gamma > 0$ and a sequence (H_ℓ) , $H_\ell \rightarrow \infty$ as $\ell \rightarrow \infty$, such that for each $\ell \geq 1$, we have

$$\overline{\lim}_{N \rightarrow \infty} \frac{1}{\log N} \sum_{1 \leq n \leq N} \frac{1}{n} \left\| \frac{1}{H_\ell} \sum_{1 \leq h \leq H_\ell} S_{n+h} f \right\|_2 \geq \gamma.$$

We write this in the form

$$\overline{\lim}_{N \rightarrow \infty} \frac{1}{\log N} \frac{1}{H_\ell} \sum_{r=0}^{H_\ell-1} \sum_{n \leq N, n \equiv r [H_\ell]} \frac{1}{n} \left\| \sum_{1 \leq h \leq H_\ell} S_{n+h} f \right\|_2 \geq \gamma.$$

Then there exists r_ℓ , $0 \leq r_\ell < H_\ell$, such that

$$\overline{\lim}_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N, n \equiv r_\ell [H_\ell]} \frac{1}{n} \left\| \sum_{1 \leq h \leq H_\ell} S_{n+h} f \right\|_2 \geq \gamma.$$

Using

$$\frac{1}{n} = \frac{1}{n+h} + \frac{h}{n(n+h)}, \quad \lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N, n \equiv r_\ell [H_\ell]} \frac{H_\ell^2}{n^2} = 0,$$

we obtain

$$\overline{\lim}_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N, n \equiv r_\ell [H_\ell]} \left\| \sum_{1 \leq h \leq H_\ell} \frac{1}{n+h} S_{n+h} f \right\|_2 \geq \gamma,$$

or

$$\overline{\lim}_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N, n \equiv r_\ell [H_\ell]} \left\| \sum_{n < m \leq n+H_\ell} \frac{1}{m} S_m f \right\|_2 \geq \gamma.$$

Rewrite this inequality in the form

$$\overline{\lim}_{N \rightarrow \infty} \frac{1}{\log N} \sum_{k=1}^{[N/H_\ell]} \left\| \sum_{kH_\ell + r_\ell < m \leq (k+1)H_\ell + r_\ell} \frac{1}{m} S_m f \right\|_2 \geq \gamma,$$

and take, for a fixed ℓ , the sequence $(b_{k,\ell})_{k \geq 1}$ defined by $b_{k,\ell} := kH_\ell + r_\ell + 1$. Setting $K_N := [N/H_\ell]$ for each N , we have $\log b_{K_N+1,\ell} / \log N \xrightarrow{N \rightarrow \infty} 1$, hence

$$\limsup_{K \rightarrow \infty} \frac{1}{\log b_{K+1,\ell}} \sum_{k \leq K} \left\| \sum_{b_{k,\ell} \leq m < b_{k+1,\ell}} \frac{1}{m} S_m f \right\|_2 \geq \gamma.$$

We can now apply the Diagonalization Lemma 3.3 with the sequences

$$g_{n,m} := \left\| \sum_{n \leq j < m} \frac{1}{j} S_j f \right\|_2.$$

We obtain that there exists a sequence $(b_k)_{k \geq 1}$ with $0 < b_{k+1} - b_k \xrightarrow{k \rightarrow \infty} \infty$, such that

$$\overline{\lim}_{K \rightarrow \infty} \frac{1}{\log b_{K+1}} \sum_{k \leq K} \left\| \sum_{b_k \leq m < b_{k+1}} \frac{1}{m} S_m f \right\|_2 \geq \gamma/2.$$

Hence (3.1) is not satisfied □

Proof of Theorem 3.1. By Proposition 3.2, we obtain that if we set

$$R_f(H) := \limsup_{N \rightarrow \infty} \frac{1}{\log N} \sum_{1 \leq n \leq N} \frac{1}{n} \left\| \frac{1}{H} \sum_{1 \leq h \leq H} S_{n+h} f \right\|_2,$$

then $\lim_{H \rightarrow \infty} R_f(H) = 0$, so the result follows from Theorem 2.1 with $\phi(s) = s$. □

Proof of Theorem 1.1. We apply Theorem 3.1 to $\mathcal{B}_1 = \mathcal{B}_2 = C(X)$ and $S_k(f) := \mu(k)f \circ T^k$. □

4. PROOF OF COROLLARY 1.2 AND RELATED RESULTS

Recall that a point y in a topological dynamical system (Y, S) is *quasi-generic* for some measure ν if, for some subsequence (N_k) of integers and all $f \in C(Y)$, we have

$$\frac{1}{N_k} \sum_{1 \leq n \leq N_k} f(S^n y) \xrightarrow{k \rightarrow \infty} \int_Y f d\nu.$$

Likewise, we say that y is *logarithmically quasi-generic* for ν if, for some subsequence (N_k) of integers and for all $f \in C(Y)$, we have

$$\frac{1}{\log N_k} \sum_{1 \leq n \leq N_k} \frac{1}{n} f(S^n y) \xrightarrow{k \rightarrow \infty} \int_Y f d\nu.$$

Observe that any measure for which y is logarithmically quasi-generic is S -invariant.

We will use here the following result from [10] (see the remark after Theorem 1.3 therein).

Theorem 4.1 (Frantzikinakis and Host). *Let (Y, S) be a topological dynamical system, and let $y \in Y$. Assume that, for any measure ν for which y is logarithmically quasi-generic, the system (Y, ν, S) has zero entropy and countably many ergodic components. Then for any $g \in C(Y)$, we have*

$$(4.1) \quad \lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{1 \leq n \leq N} \frac{1}{n} g(S^n y) \mu(n) = 0.$$

Proof of Corollary 1.2. Let us consider a dynamical system (X, T) with zero topological entropy, and such that $M^e(X, T)$ is countable. In view of Theorem 1.1, all we need to prove is that (X, T) satisfies the logarithmic strong MOMO property. That is, we fix an increasing sequence $(b_k) \subset \mathbb{N}$ with $b_{k+1} - b_k \rightarrow \infty$ (and we assume without loss of generality that $b_1 = 1$), a sequence $(x_k) \subset X$ and $f \in C(X)$, and we have to show the following convergence

$$(4.2) \quad \frac{1}{\log b_{K+1}} \sum_{k \leq K} \left| \sum_{b_k \leq n < b_{k+1}} \frac{1}{n} f(T^n x_k) \mu(n) \right| \xrightarrow{K \rightarrow \infty} 0.$$

According to [3, Lemma 18], it is sufficient to show that

$$(4.3) \quad \frac{1}{\log b_{K+1}} \sum_{k \leq K} e_k \sum_{b_k \leq n < b_{k+1}} \frac{1}{n} f(T^n x_k) \mu(n) \xrightarrow{K \rightarrow \infty} 0,$$

where $e_k \in \Sigma_3 := \{e^{2\pi i j/3} : j = 0, 1, 2\}$ is chosen so that the product

$$e_k \sum_{b_k \leq n < b_{k+1}} \frac{1}{n} f(T^n x_k) \mu(n)$$

belongs to the closed cone $\{0\} \cup \{z \in \mathbb{C} : \arg(z) \in [-\pi/3, \pi/3]\}$.

In order to show (4.3), we consider the space $Y := (X \times \Sigma_3)^{\mathbb{N}}$ with the shift S , and in this system the point $y = (y_n)_{n \in \mathbb{N}}$ defined by

$$y_n := (T^n x_k, e_k) \text{ if } b_k \leq n < b_{k+1} \quad (k \geq 1).$$

Let ν be a measure for which y is logarithmically quasi-generic. The same argument as in [3] (see the proof of $(P2 \Rightarrow P3)$) shows that ν must be concentrated on the set of sequences of the form

$$((x, a), (Tx, a), (T^2x, a), \dots) \quad (x \in X, a \in \Sigma_3).$$

Now, let us consider an ergodic component ρ of ν . The marginal of ρ on the first coordinate x must be an ergodic T -invariant measure on X . By assumption, there are only countably many of them, and all of them give rise to zero-entropy systems. The marginal of ρ on the second coordinate a is one of the three Dirac measures $\delta_1, \delta_{e^{i2\pi/3}}, \delta_{e^{i4\pi/3}}$. Moreover these two marginals must be independent by the disjointness of ergodic systems with the identity, thus, these two marginals completely determine ρ . Hence, we see that there can be only countably many possible ergodic components of ν , and all of them have zero entropy. Thus y satisfies the assumptions of Theorem 4.1, and we have (4.1) for each $g \in C(Y)$. In particular, if we take the continuous function g defined by $g(z) := a_0 f(z_0)$ for each $z = ((z_0, a_0), (z_1, a_1), \dots) \in Y$, we obtain (4.3). \square

Remark 4.2. We can characterize uniform convergence for Möbius orthogonality in terms of a MOMO type convergence. Indeed:

The uniform convergence in Möbius orthogonality (1.1) holds if and only if for all (b_k) satisfying $b_k/b_{k+1} \rightarrow 0$, we have

$$\frac{1}{b_{k+1}} \left\| \sum_{b_k \leq n < b_{k+1}} \mu(n) f \circ T^n \right\|_{C(X)} \xrightarrow{k \rightarrow \infty} 0.$$

To see this equivalence, it is sufficient to note that for each $x \in X$,

$$\begin{aligned} \frac{1}{b_{k+1}} \sum_{b_k \leq n < b_{k+1}} f(T^n x) \mu(n) &= \\ \frac{1}{b_{k+1}} \sum_{1 \leq n < b_{k+1}} f(T^n x) \mu(n) - \frac{b_k}{b_{k+1}} \frac{1}{b_k} \sum_{1 \leq n < b_k} f(T^n x) \mu(n). \end{aligned}$$

Remark also that the same arguments work for logarithmic averages (replacing $b_k/b_{k+1} \rightarrow 0$ with $\log b_k / \log b_{k+1} \rightarrow 0$).

5. MISCELLANEA

5.1. Ergodic measures. In this section, we show that Tao's approach persists, if we consider the main observation from [12].

Let (X, T) be a dynamical system. Given $x \in X$ and $n \in \mathbb{N}$, we write $\delta_{T^n(x)}$ for the Dirac measure concentrated at the point $T^n(x)$. Let

$$\mathcal{E}(x, N) := \frac{1}{N} \sum_{1 \leq n \leq N} \delta_{T^n(x)}, \quad \mathcal{E}^{\log}(x, N) := \frac{1}{\log N} \sum_{1 \leq n \leq N} \frac{1}{n} \delta_{T^n(x)}.$$

We consider here the convergence of these empirical measures in the weak* topology. Note that any accumulation point of the above sequences is always a T -invariant probability measure on X .

Theorem 5.1. *Suppose that for some $x \in X$ and some subsequence $(N_s)_{s \geq 1}$ of natural numbers, we have*

$$(5.1) \quad \lim_{s \rightarrow \infty} \mathcal{E}^{\log}(x, N_s) = \kappa, \quad \text{where } \kappa \text{ is ergodic.}$$

Then there exists a set \mathcal{N} of natural numbers with the property

$$(5.2) \quad \lim_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{\mathcal{N} \ni N \leq N_s} \frac{1}{N} = 1,$$

such that

$$(5.3) \quad \lim_{N \rightarrow \infty, N \in \mathcal{N}} \mathcal{E}(x, N) = \kappa.$$

Proof. The condition (5.1) means that for any $f \in C(X)$ we have

$$(5.4) \quad \lim_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{n=1}^{N_s} \frac{1}{n} f(T^n u) = \int_X f d\kappa.$$

Let $f \in C(X)$ be fixed, and set $Sf := \int_X f d\kappa$. For $H \in \mathbb{N}$, we consider the limiting of the second moment

$$R_f(H) := \lim_{s \rightarrow \infty} \frac{1}{\log N_s} \sum_{1 \leq n \leq N_s} \frac{1}{n} \left| \frac{1}{H} \sum_{m=1}^H f(T^{n+m} x) - Sf \right|^2.$$

The limit does exist by (5.1), as the internal is given by a continuous function sum sampled at x . So, by condition (5.4), we have

$$(5.5) \quad R_f(H) = \int_X \left| \frac{1}{H} \sum_{m=1}^H f(T^m x) - Sf \right|^2 d\kappa(x).$$

Hence, directly by the von Neumann ergodic theorem, and using the ergodicity of κ , we obtain $\lim_{H \rightarrow \infty} R_f(H) = 0$. Take now $\mathcal{B}_1 = C(X)$, $\mathcal{B}_2 = \mathbb{C}$ with the sequence of functionals $S_k f := f(T^k x) - Sf$, $k \in \mathbb{N}$, and we obtain statement (5.3) by Theorem 2.1. \square

5.2. Davenport-Erdős theorem. Davenport-Erdős theorem [7] is the fact that, given $\mathcal{B} \subset \mathbb{N}$, the \mathcal{B} -free set $\mathcal{F}_{\mathcal{B}}$, i.e. the set of those numbers that have no divisor in \mathcal{B} , has logarithmic density and, moreover, $\delta(\mathcal{F}_{\mathcal{B}}) = \bar{d}(\mathcal{F}_{\mathcal{B}})$. In fact, see [8], the point $\mathbb{1}_{\mathcal{F}_{\mathcal{B}}}$ is **logarithmically generic** for the relevant Mirsky measure which is ergodic. Hence, by Theorem 5.1, we obtain that the upper asymptotic density $\bar{d}(\mathcal{F}_{\mathcal{B}})$ is obtained along a subsequence of logarithmic density 1. We can however obtain this result in an elementary way. Indeed, for a subset $A \subset \mathbb{N}$ and $N \in \mathbb{N}$, set $d_N(A) := \frac{1}{N} \sum_{1 \leq n \leq N} \mathbb{1}_A(n)$ and $d_N^{\log}(A) := \frac{1}{\log N} \sum_{1 \leq n \leq N} \frac{1}{n} \mathbb{1}_A(n)$. More generally, given $a = (a_n)_{n \in \mathbb{N}}$ a sequence of real numbers, set $d_N(a) := \frac{1}{N} \sum_{1 \leq n \leq N} a_n$ and $d_N^{\log}(a) := \frac{1}{\log N} \sum_{1 \leq n \leq N} \frac{a_n}{n}$.

Proposition 5.2. *Let $a = (a_n)_{n \in \mathbb{N}}$ be a bounded sequence of real numbers, and let $\ell := \limsup_{N \rightarrow \infty} d_N(a)$. Assume that $\lim_{N \rightarrow \infty} d_N^{\log}(a) = \ell$. Then there exists $B \subset \mathbb{N}$ with $\lim_{N \rightarrow \infty} d_N^{\log}(B) = 1$ such that $\lim_{B \ni N \rightarrow \infty} d_N(a) = \ell$.*

Proof. Step 1 For any $\varepsilon > 0$, set $B_\varepsilon := \{N \in \mathbb{N} : d_N(a) > \ell - \varepsilon\}$. Then $\lim_{N \rightarrow \infty} d_N^{\log}(B_\varepsilon) = 1$. Indeed, let us introduce the sequence $b = (b_n)_{n \in \mathbb{N}}$ defined by $b_1 := 0$, and for each $n \geq 2$, $b_n := d_{n-1}(a)$. The Abel summation formula yields

$$d_N^{\log}(a) = \frac{d_N(a)}{\log N} + d_N^{\log}(b).$$

By assumption, we thus have $\lim_{N \rightarrow \infty} d_N^{\log}(b) = \ell$. Let

$$\widetilde{B}_\varepsilon := \{N \in \mathbb{N} : b_N > \ell - \varepsilon\} (= B_\varepsilon + 1) \text{ and } \widetilde{A}_\varepsilon := \mathbb{N} \setminus \widetilde{B}_\varepsilon = \{N \in \mathbb{N} : b_N \leq \ell - \varepsilon\}.$$

In the computation of $d_N^{\log}(b)$, the contribution of $\widetilde{A}_\varepsilon$ is bounded above by

$$(\ell - \varepsilon) d_N^{\log}(\widetilde{A}_\varepsilon) = (\ell - \varepsilon) \left(1 - d_N^{\log}(\widetilde{B}_\varepsilon)\right).$$

On the other hand, using the fact that $\limsup_{N \rightarrow \infty} b_N = \ell$, the contribution of $\widetilde{B}_\varepsilon$ to $d_N^{\log}(b)$ is bounded above by $\ell d_N^{\log}(\widetilde{B}_\varepsilon) + o(1)$. Therefore, we have for each $N \in \mathbb{N}$

$$\begin{aligned} d_N^{\log}(b) &\leq (\ell - \varepsilon) \left(1 - d_N^{\log}(\widetilde{B}_\varepsilon)\right) + \ell d_N^{\log}(\widetilde{B}_\varepsilon) + o(1) \\ &= \varepsilon d_N^{\log}(\widetilde{B}_\varepsilon) + \ell - \varepsilon + o(1). \end{aligned}$$

But we know that $\lim_{N \rightarrow \infty} d_N^{\log}(b) = \ell$, and it follows that $\lim_{N \rightarrow \infty} d_N^{\log}(\widetilde{B}_\varepsilon) = 1$.

Finally, since $\widetilde{B}_\varepsilon = B_\varepsilon + 1$, we also get $\lim_{N \rightarrow \infty} d_N^{\log}(B_\varepsilon) = 1$.

Step 2 We construct the announced set B as follows. First we fix a decreasing sequence (ε_k) of positive numbers going to 0 as $k \rightarrow \infty$. By Step 1, we know that $d_N^{\log}(B_{\varepsilon_k}) \xrightarrow{N \rightarrow \infty} 1$. Then we construct a strictly increasing sequence (N_k) of integers such that $\forall k, \forall N \geq N_k, d_N^{\log}(B_{\varepsilon_k}) > 1 - \varepsilon_k$. Finally we define B by $B \cap \{1, \dots, N_1 - 1\} := \{1, \dots, N_1 - 1\}$, and for each $k \geq 1$, $B \cap \{N_k, \dots, N_{k+1} - 1\} := B_{\varepsilon_k} \cap \{N_k, \dots, N_{k+1} - 1\}$. \square

5.3. Deriving a density version of the PNT from the logarithmic Chowla conjecture of order 2.

Lemma 5.3. *Assume that $A \subset \mathbb{N}$ with $\delta(A) = 1$. For each $m \in \mathbb{N}$ set*

$$(5.6) \quad A_m := \{n \in \mathbb{N} : [n/m] \in A\}.$$

Then $\delta(A_m) = 1$ for all $m \in \mathbb{N}$.

Proof. Let $m \geq 2$ be fixed. Note that if $n \in A$, then

$$(5.7) \quad mn + j \in A_m, \quad j = 0, 1, \dots, m-1,$$

and

$$(5.8) \quad \sum_{j=0}^{m-1} \frac{1}{mn+j} \geq \int_{mn}^{m(n+1)} \frac{dt}{t} = \log(1 + 1/n) \geq \frac{1}{n} - \frac{1}{2n^2}.$$

Given $\epsilon \in (0, 1)$, we have $\frac{1}{\log N} \sum_{n \in A, n \leq N} \frac{1}{n} \geq 1 - \epsilon/2$ for $N \geq N_\epsilon$. Setting $c := \sum_{k=1}^{\infty} \frac{1}{k^2} = \frac{\pi^2}{6}$, we may assume that

$$\frac{c}{2 \log N_\epsilon} \leq \epsilon/2 \quad \text{and} \quad \frac{\log N}{\log mN} \geq 1 - \epsilon \quad \text{for all } N \geq N_\epsilon.$$

It follows that for any $N \geq N_\epsilon$, by (5.8) and (5.7), we have

$$\begin{aligned} 1 - \epsilon/2 &\leq \frac{1}{\log N} \sum_{n \in A, n \leq N} \frac{1}{n} \\ &\leq \frac{1}{\log N} \sum_{n \in A, n \leq N} \left(\sum_{j=0}^{m-1} \frac{1}{mn+j} \right) + \frac{1}{2 \log N} \sum_{n \in A, n \leq N} \frac{1}{n^2} \\ &\leq \frac{1}{\log N} \sum_{k \in A_m, k \leq m(N+1)} \frac{1}{k} + \epsilon/2. \end{aligned}$$

Therefore, for all $N \geq N_\epsilon$, we have

$$1 \geq \frac{1}{\log mN} \sum_{k \in A_m, k \leq mN} \frac{1}{k} = \frac{\log N}{\log mN} \frac{1}{\log N} \sum_{k \in A_m, k \leq mN} \frac{1}{k} \geq (1 - \epsilon)^2.$$

Letting $\epsilon \rightarrow 0$, we obtain $\lim_{N \rightarrow \infty} \frac{1}{\log mN} \sum_{k \in A_m, k \leq mN} \frac{1}{k} = 1$, and then $\delta(A_m) = 1$. \square

By Lemmas 5.3 and 2.4 we obtain the following:

Lemma 5.4. *Let $A \subset \mathbb{N}$ with $\delta(A) = 1$. Then there exists $\tilde{A} \subset \mathbb{N}$ with $\delta(\tilde{A}) = 1$, such that for any $m \in \mathbb{N}$ there exists N_m satisfying*

$$(5.9) \quad \tilde{A} \cap \{n : n \geq N_m\} \subset \bigcap_{k=1}^m A_k$$

(sets A_k are defined by (5.6)).

Given $u : \mathbb{N} \rightarrow \mathbb{C}$, let $U(x) := \sum_{n \leq x} u(n)$, for $x \geq 0$, denote the corresponding summation function.

Lemma 5.5. *Let $A \subset \mathbb{N}$ with $\delta(A) = 1$, and let $u : \mathbb{N} \rightarrow \mathbb{C}$ such that*

$$\frac{|U(n)|}{n} \xrightarrow{A \ni n \rightarrow \infty} 0.$$

Then there exists $\tilde{A} \subset A$, $\delta(\tilde{A}) = 1$ such that, for each $a \geq 1$ and $\varepsilon > 0$, we can find $X = X(a, \varepsilon) > 1$ for which

$$\forall x \geq X, [x] \in \tilde{A} \implies \sum_{n \leq a} |U(x/n)| \leq \varepsilon x.$$

Proof. We have a set \tilde{A} as in Lemma 5.4, so $\delta(\tilde{A}) = 1$. Let $a \geq 1$ be fixed. Denote $C := \sum_{n \leq a} \frac{1}{n}$ and choose $K \geq 1$ so that

$$(5.10) \quad A \ni [x] \geq K \implies \frac{|U(x)|}{x} \leq \frac{|U([x])|}{[x]} \leq \varepsilon/C.$$

Taking $m = [a]$ and using Lemma 5.4, we choose N_m such that (5.9) holds, i.e. $[n], [n/2], \dots, [n/m] \in A$ whenever $n \in \tilde{A}$ and $n \geq N_m$. Then for $x \geq (m+1) \max\{N_m, K\}$ with $[x] \in \tilde{A}$, by (5.10), we have $\sum_{n \leq a} |U(x/n)| \leq \sum_{n \leq a} \frac{x}{n} \frac{\varepsilon}{C} = x\varepsilon$. \square

Proposition 5.6. *The statement logarithmic Chowla conjecture (for μ) holds⁶ for auto-correlations of length 2 implies that there exists a sequence $A \subset \mathbb{N}$, $\delta(A) = 1$, such that $\sum_{n \leq x} \Lambda(n) = x + o(x)$ for $A \ni x \rightarrow \infty$.*

Proof. By Theorem 2.1⁷ (or directly by Tao's proof in [18]), we obtain that $\frac{1}{N} \sum_{n \leq N} \mu(n) \rightarrow 0$ when $A \ni N \rightarrow \infty$, where $\delta(A) = 1$.

Following [4], we repeat the proof that $M(\mu) := \lim_{N \rightarrow \infty} \sum_{n \leq N} \mu(n) = 0$ implies PNT. We have

$$\sum_{n \leq x} \Lambda(n) = x - \sum_{q, d; qd \leq x} \mu(d) f(q) + O(1)$$

for some arithmetic function f . All we need to show is that $\sum_{q, d; qd \leq x} \mu(d) f(q) = o(x)$. Using summation by parts, one arrives at

$$(5.11) \quad \sum_{q, d; qd \leq x} \mu(d) f(q) = \sum_{n \leq b} \mu(n) F(x/n) + \sum_{n \leq a} f(n) M(x/n) - F(a) M(b),$$

where $ab = x$. Moreover, $F(x) = B \cdot \sqrt{x}$ for a constant $B > 0$. It follows that the first summand in (5.11) is bounded by $B_1 x / \sqrt{a}$ for a constant $B_1 > B$. Indeed,

$$\left| \sum_{n \leq b} \mu(n) F(x/n) \right| \leq B \sqrt{x} \sum_{n \leq b} \frac{1}{\sqrt{n}} \leq B_1 \sqrt{b} \sqrt{x} = B_1 \frac{x}{\sqrt{a}}.$$

We fix $\varepsilon > 0$ and choose $a \geq 1$, so that $B_1 / \sqrt{a} < \varepsilon$ which yields the first summand $< \varepsilon x$ for all $x \geq 1$. To majorate the second summand, we use

$$\left| \sum_{n \leq a} f(n) M(x/n) \right| \leq F_a \sum_{n \leq a} |M(x/n)|, \quad F_a := \max\{|f(n)| : n \leq a\},$$

and Lemma 5.5 (with $u = \mu$). Finally, the third summand is majorated in the same way as in [4]. \square

Acknowledgments. Research of the first and second authors supported by Narodowe Centrum Nauki grant UMO-2019/33/B/ST1/00364.

⁶Proved by Tao in [16].

⁷Note that if (c_n) is a bounded sequence of complex numbers with $\lim_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N} \frac{1}{n} c_n \overline{c_{n+h}} = 0$ for each $h \neq 0$ then for

$$R_H((c_n)) := \limsup_{N \rightarrow \infty} \frac{1}{\log N} \sum_{n \leq N} \frac{1}{n} \left| \frac{1}{H} \sum_{h \leq H} c_{n+h} \right|^2$$

we have $\lim_{H \rightarrow \infty} R_H((c_n)) = 0$.

REFERENCES

- [1] E. H. El Abdalaoui, M. Lemańczyk and T. de la Rue, *Automorphisms with quasi-discrete spectrum, multiplicative functions and average orthogonality along short intervals*, International Math. Res. Notices **14** (2017), 4350–4368.
- [2] E. H. El Abdalaoui, J. Kułaga-Przymus, M. Lemańczyk, and T. de la Rue, *The Chowla and the Sarnak conjectures from ergodic theory point of view*, Discrete Contin. Dyn. Syst. **37** (2017), 2899–2944.
- [3] E. H. El Abdalaoui, J. Kułaga-Przymus, M. Lemańczyk, T. de la Rue, *Möbius disjointness for models of an ergodic system and beyond*, Israel J. Math. **228** (2018), 707–751.
- [4] H. Apostol, *Introduction to Analytic Number Theory*, Undergraduate Texts in Mathematics, Springer, 2010.
- [5] J. Bourgain, P. Sarnak and T. Ziegler, *Disjointness of Möbius from horocycle flows*, From Fourier analysis and number theory to Radon transforms and geometry, Dev. Math., vol. 28, Springer, New York, 2013, 67–83.
- [6] F. Cellarosi and Ya. G. Sinai, *Ergodic properties of square-free numbers*, J. European Math. Soc. **15** (2013), 1343–1374.
- [7] H. Davenport and P. Erdős, *On sequences of positive integers*, Acta Arithmetica **2** (1936), 147–151.
- [8] A. Dymek, S. Kasjan, J. Kułaga-Przymus and M. Lemańczyk, *B-free sets and dynamics*, Trans. Amer. Math. Soc. **370** (2018), 5425–5489.
- [9] S. Ferenczi, J. Kułaga-Przymus and M. Lemańczyk, *Sarnak’s conjecture, what’s new*, in: Ergodic Theory and Dynamical Systems in their Interactions with Arithmetics and Combinatorics, S. Ferenczi, J. Kułaga-Przymus, M. Lemańczyk (eds), Lecture Notes in Math. vol. 2213, Springer, Cham, 2018, pp. 163–235.
- [10] N. Frantzikinakis and B. Host, *The logarithmic Sarnak conjecture for ergodic weights*, Annals Math. **187** (2018), 869–931.
- [11] N. Frantzikinakis and B. Host, *Furstenberg systems of bounded multiplicative functions and applications*, to appear in International Math. Res. Notices, arXiv:1804.08556.
- [12] A. Gomilko, D. Kwietniak and M. Lemańczyk, *Sarnak’s conjecture implies the Chowla conjecture along a subsequence*, in: Ergodic Theory and Dynamical Systems in their Interactions with Arithmetics and Combinatorics, S. Ferenczi, J. Kułaga-Przymus, M. Lemańczyk (eds), Lecture Notes in Math. vol. 2213, Springer, Cham, 2018, pp. 237–247.
- [13] A. Kanigowski, M. Lemańczyk and C. Ulcigrai, *On disjointness of some parabolic flows*, Inventiones Math. **221** (2020), 1–111.
- [14] P. Sarnak, *Three lectures on the Möbius function, randomness and dynamics*, <http://publications.ias.edu/sarnak/>.
- [15] T. Tao, <https://terrytao.wordpress.com/2012/10/14/the-chowla-conjecture-and-the-sarnak-conjecture/>
- [16] T. Tao, *The logarithmically averaged Chowla and Elliot conjectures for two-point correlations*, Forum Math. Pi **4** (2016).
- [17] T. Tao, *Equivalence of the logarithmically averaged Chowla and Sarnak conjectures*, Number Theory – Diophantine Problems, Uniform Distribution and Applications: Festschrift in Honour of Robert F. Tichy’s 60th Birthday (C. Elsholtz and P. Grabner, eds.), Springer International Publishing, Cham, 2017, 391–421.
- [18] T. Tao, <https://terrytao.wordpress.com/2017/10/20/the-logarithmically-averaged-and-non-logarithmically-averaged-chowla-conjectures/>
- [19] T. Tao, J. Teräväinen, *Odd order cases of the logarithmically averaged Chowla conjecture*, J. Théor. Nombres de Bordeaux **30** (2018), 997–1015.
- [20] T. Tao, J. Teräväinen, *The structure of logarithmically averaged correlations of multiplicative functions, with applications to the Chowla and Elliot conjectures*, Duke Math. J. **168** (2019), 1977–2027.

*Manuscript received June 3 2019**revised August 4 2020*

ALEXANDER GOMILKO

Faculty of Mathematics and Computer Science, Nicolaus Copernicus University, Chopin street
12/18, 87-100 Toruń, Poland*E-mail address:* `gomilko@mat.umk.pl`

MARIUSZ LEMAŃCZYK

Faculty of Mathematics and Computer Science, Nicolaus Copernicus University, Chopin street
12/18, 87-100 Toruń, Poland*E-mail address:* `mlem@mat.umk.pl`

THIERRY DE LA RUE

Laboratoire de Mathématiques Raphaël Salem, Université de Rouen Normandie, CNRS – Avenue
de l'Université – 76801 Saint Etienne du Rouvray, France*E-mail address:* `Thierry.de-la-Rue@univ-rouen.fr`